

09/889756 12310
JC14 Rec'd PCT/PTO 20 JUL 2001
09/889756 PATENT
DOCKET NO. BM45353


IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

BOX PCT/DO-EO

Commissioner for Patents
Washington, D.C. 20231

"Express Mail" mailing label number **EL 742 362 336 US**

I hereby certify that this paper is being deposited with the United States Postal Service "Express Mail Post Office to Addressee" under 37 CFR, § 1.10 on the date indicated above and addressed to: Commissioner of Patents, Washington, DC 20231, on July 20, 2001.


Colleen Hanagan

NATIONAL STAGE APPLICATION TRANSMITTAL LETTER
APPLICATION FILING UNDER 35 U.S.C. § 371

Transmitted herewith for filing is the patent application of:

Inventor(s)/Applicant(s): Thonnard, Joelle
International Application No.: PCT/EP00/00425
International Published Appln. No.: WO 00/43517
International Filing Date: 19 January 2000
Priority Filing Dates: 22 January 1999 & 29 January 1999
Thirty Month Date: 22 July 2001
Title: "BASB055 POLYNUCLEOTIDE AND POLYPEPTIDE FROM NEISSERIA MENINGITIDIS, USES THEREOF"

1. **THIS NEW APPLICATION IS A NATIONAL STAGE APPLICATION UNDER PCT, CHAPTER II WITH A REQUEST FOR EXAMINATION WITHOUT DELAY TO THE UNITED STATES DESIGNATED/ELECTED OFFICE (DO/EO/US).**

- ☒ This is a **FIRST** submission of items concerning a filing under 35 U.S.C. § 371;
☐ This is a **SECOND** or subsequent submission of items concerning a filing under 35 U.S.C. § 371.

2. ☒ This is an express request to begin national examination procedures (35 U.S.C. 371(f)) at any time rather than delay examination until the expiration of the applicable time limit 35 U.S.C. 371(b) and PCT Articles 22 and 39(1).

3. A proper Demand for International Preliminary Examination was made by the 19TH month from the earliest claimed priority date.

4. Enclosed items are required for filing under 37 CFR § 1.53(b) and § 1.494(b) or § 1.495(b):

- ☒ One copy of International Publication No. WO 00/43517
- (a) ☒ is transmitted herewith (**required only if not transmitted by the International Bureau**)
- (b) ☐ has been transmitted by the International Bureau
- (c) ☐ is not required, as the application was filed in the United States Receiving Office (RO/US)

Fees

☒ The basic national fee set forth in 37 CFR § 1.482 - International Preliminary Examination Fee not paid to USPTO but International Search Report prepared by the EPO or JPO - **\$860.00**

- ☒ Claims in Excess of 20 (4 @ \$18.00)
- ☐ Independent Claims in Excess of 3 (@ \$80.00)

5. Further enclosed are:

- ☒ One copy of International Preliminary Examination Report.
- ☒ One copy of International Search Report.
- ☐ One copy of Written Opinion.
- ☐ One copy of PCT Request as filed.
- ☐ One copy of Chapter II Demand as filed.

6. ☐ A translation of the International Application into English (35 U.S.C. § 371(c)(2))

7. ☐ Amendments to the claims of the International Application under PCT Article 19 (35 U.S.C. § 371(c)(3))

8. ☐ A translation of the amendments to the claims under PCT Article 19 (35 U.S.C. § 371(c)(3))

9. Still additional papers enclosed:

- ☐ Assignment with Assignment Recordation Form Cover Sheet
- ☐ Verification Statement Claiming Small Entity Status
- ☐ Declaration or oath is enclosed executed by the inventor
- ☐ An Information Disclosure Statement under 37 CFR § 1.97 and § 1.98
- ☒ Return Acknowledgment Postcard

10. Nucleotide and/or Amino Acid Sequence Submission (if applicable, all necessary)
- A. Enclosed are:
- (a) ☐ Computer Readable Copy of the Sequence Listing
- (b) ☐ Paper Copy (identical to Computer Readable Copy) of the Sequence Listing
- B. ☐ Enclosed is a paper copy of the Sequence Listing. This paper copy and a Computer Readable Form thereof are identical with the Computer Readable Form in another application of the Applicant which is fully identified as follows:

U.S. Application No.: @@

Filed: @@

Attorney Docket No.: @@

which is believed to comply with the rules set forth in 37 CFR § 1.821 et. seq. Applicants requests pursuant to 37 CFR § 1.821(e) that this Computer Readable Form be used in the present application. **Please TRANSFER the sequence listing from the parent to this application.**

- C. ☐ Statement under 37 CFR § 1.821(f): **The information recorded in computer readable form is identical to the written Sequence Listing.**
- D. ☐ Statement under 37 CFR § 1.821(g) (required when Sequence Listing not submitted at the time of filing under 35 U.S.C. § 111(a)) or 37 CFR § 1.821(f) (required when Sequence Listing not submitted at the time of filing under the Patent Cooperation Treaty): **The submission of the Sequence Listing includes no new matter.**
- E. ☐ Amendment: Please enter the Sequence Listing into the application.

11. **Preliminary Amendment**

Prior to calculation of fees, kindly enter:

- ☒ Preliminary Amendment submitted herewith
- ☐ do not enter Preliminary Amendment

09/889756

09/889756

PATENT

DOCKET NO. BM45353

JC18 Rec'd PCT/PTO 20 JUL 2001

12. The correspondence address for this application is the Customer No. provided below:

Insert Bar Code Label Here:



25308

PATENT, TRADEMARK OFFICE

13. Fee payment being made at this time is enclosed:

* Basic filing fee (\$860.00)	860.00
* Claims in Excess of 20	72.00
* (4 @ \$18.00)	
Independent Claims in Excess of 3	00.00
(@ \$80.00)	
* Total Fees enclosed:	<u>\$932.00</u>

14. The Commissioner is hereby authorized to charge any fees which may be required, or credit any overpayment to Account No. 50-0258. This letter is filed in duplicate for accounting purposes.

Respectfully submitted,

E. A. Meade

Eric A. Meade

Registration No. 42,876

for

Allen Bloom

Registration No. 29,135

Attorney for Applicant

DECHERT
Princeton Pike Corporate Center
PO Box 5218
Princeton, New Jersey 08543-5218
Allen Bloom (609) 620-3214
Eric A. Meade (609) 620-3248
Fax: (609) 620-3259
Attn.: Allen Bloom, Esq.
(609) 620-3214

C:\WIN98\TEMP.2H2401\DOC

Docket No: BM45353

Rec'd PCT/PTO 31 DEC 2001
09/889756 #3

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re Application of	:	International Application No.: PCT/EP00/00425
Thonnard	:	I.A. Filing Date: 19 January 2000
Serial No.: 09/889,756	:	Priority Dates: 22 January 1999 and
	:	29 January 1999
	:	
For: BASB055 Polynucleotide and	:	
Polypeptide From <i>Neisseria</i>	:	
<i>Meningitidis</i> , Uses Thereof	:	

Commissioner for Patents
Box PCT
Washington, DC 20231

**RESPONSE TO NOTIFICATION TO COMPLY WITH REQUIREMENTS
FOR PATENT APPLICATIONS CONTAINING NUCLEOTIDE SEQUENCE
AND/OR AMINO ACID SEQUENCE DISCLOSURES**

Sir:


This is in reply to the communication mailed 20 August 2001 from the Patent Office, entitled "Notice to Comply with Requirements for Patent Applications Containing Nucleotide Sequence and/or Amino Acid Sequence Disclosures." A copy of the Notice is enclosed.

Submitted herewith is the "Sequence Listing" in computer readable form in accordance with the requirements of 37 CFR §§ 1.821-1.825.

STATEMENT

Applicant hereby states that the enclosed CRF diskette submitted herewith is identical to the paper copy of the "Sequence Listing" contained in the copy of International Publication No. WO 00/43517 which was submitted on July 20, 2001 for filing with the US Patent Office and contains no new matter.

Respectfully submitted,



Arthur E. Jackson
Registration No. 34,354
Attorney for Applicant

DECHERT
4000 Bell Atlantic Tower
1717 Arch Street
Philadelphia, PA 19103-2793
Fax: (609) 620-3259
Attn: Eric A. Meade
(609) 620-3248

Docket No. BM45353
Serial No. 09/889,756

CERTIFICATE OF MAILING

I hereby certify that this correspondence is being caused to be deposited on the date shown below with the United States Postal Service with sufficient postage as first class mail in an envelope addressed to:
Commissioner for Patents, Box PCT, Washington, D.C. 20231.

Colleen Hanagan
Colleen Hanagan

October 19, 2021
Date

International Application No.: PCT/EP00/00425
Attorney Docket No.: BM45353

09889756 12310
09/889756
JC18 Rec'd PCT/PTO 20 JUL 2001

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

Applicant: Thonnard
Docket No.: BM45353
Serial No.: Unknown
Filed: Herewith

Group Art Unit No.: Unknown
Examiner: Unknown

For: BASB055 Polynucleotides and Polypeptides from *Neisseria Meningitidis*, Uses Thereof

PRELIMINARY AMENDMENT

Sir:

Applicant respectfully requests that this Preliminary Amendment be entered in this case before the calculation of fees and before examination of the subject application.

In the Claims:

Please delete the claims of the application as filed in the PCT and substitute therefor:

19. An isolated polypeptide comprising a member selected from the group consisting of
 - (a) an amino acid sequence matching SEQ ID NO:2;
 - (b) an immunogenic polypeptide comprising a fragment sequence of at least 15 amino acids that matches an aligned contiguous segment of SEQ ID NO:2,wherein the isolated polypeptide, when administered to a subject in a suitable composition which can include an adjuvant, or a suitable carrier coupled to the polypeptide, induces an antibody or T-cell immune response to a polypeptide having the sequence of SEQ ID NO:2.
20. An isolated polynucleotide encoding a polypeptide of Claim 19 or the full complement to the isolated polynucleotide.
21. The isolated polypeptide of claim 19, wherein the polypeptide is according to (a).
22. An isolated polynucleotide encoding a polypeptide of Claim 21 or the full complement to the isolated polynucleotide.
23. A process for expressing the polynucleotide of Claim 22 comprising transforming a host cell with an expression vector comprising the polynucleotide and culturing the host cell under conditions sufficient for expression of the polynucleotide.

International Application No.: PCT/EP00/00425
Attorney Docket No.: BM45353

24. The isolated polypeptide of claim 19, wherein the polypeptide is according to (b).
25. An isolated polynucleotide encoding a polypeptide of Claim 24 or the full complement to the isolated polynucleotide.
26. The isolated polypeptide of claim 19, wherein the immunogenic fragment of (b) comprises at least 20 amino acids.
27. The isolated polypeptide of claim 19, wherein the isolated polypeptide consists of SEQ ID NO:2.
28. An isolated polynucleotide encoding the polypeptide of Claim 28 or the full complement to the isolated polynucleotide.
29. A process for expressing the polynucleotide of Claim 28 comprising transforming a host cell with an expression vector comprising the polynucleotide and culturing the host cell under conditions sufficient for expression of the polynucleotide.
30. A fusion protein comprising the isolated polypeptide of Claim 19.
31. An isolated polynucleotide comprising the polynucleotide of SEQ ID NO:1.
32. An isolated polynucleotide segment comprising a polynucleotide sequence or the full complement of the entire length of the polynucleotide sequence, wherein the polynucleotide sequence hybridizes to the full complement of SEQ ID NO:1 minus the complement of any stop codon, wherein the hybridization conditions include incubation at 42°C in a solution comprising: 50% formamide, 5x SSC (150mM NaCl, 15mM trisodium citrate), 50 mM sodium phosphate (pH 7.6), 5x Denhardt's solution, 10% dextran sulfate, and 20 micrograms/ml denatured, sheared salmon sperm DNA, followed by washing in 0.1x SSC at 65°C; and, wherein the polynucleotide sequence is identical to SEQ ID NO:1 minus any terminal stop codon, except that, over the entire length corresponding to SEQ ID NO:1 minus any terminal

International Application No.: PCT/EP00/00425
Attorney Docket No.: BM45353

stop codon, n_n nucleotides are substituted, inserted or deleted, wherein n_n satisfies the following expression

$$n_n \leq x_n - (x_n \cdot y)$$

wherein x_n is the total number of nucleotides in SEQ ID NO:1 minus any terminal stop codon, y is at least 0.95, and wherein any non-integer product of x_n and y is rounded down to the nearest integer before subtracting the product from x_n ; and wherein the polynucleotide sequence detects *Neisseria meningitidis*.

33. An expression vector comprising the isolated polynucleotide of Claim 20.
34. A host cell transformed with the expression vector of Claim 33.
35. A vaccine comprising the polypeptide of Claim 19 and a pharmaceutically acceptable carrier.
36. The vaccine of Claim 35, wherein the vaccine comprises at least one other *Neisseria meningitidis* antigen.
37. An antibody immunospecific for the polypeptide or immunogenic fragment of Claim 19.
38. A method for inducing an immune response in a mammal comprising administration of the polypeptide of Claim 19.
39. A method of diagnosing a *Neisseria meningitidis* infection, comprising identifying a polypeptide of Claim 19, or an antibody that is immunospecific for the polypeptide, present within a biological sample from an animal suspected of having such an infection.
40. A method for inducing an immune response in a mammal comprising administration of the isolated polynucleotide of Claim 20.

[illegible]

41. A therapeutic composition useful in treating humans with *Neisseria meningitidis* comprising at least one antibody directed against the polypeptide of claim 19 and a suitable pharmaceutical carrier.

42. The isolated polypeptide of claim 19, wherein the immune response recognizes an antigen having molecular weight of about 50 kDa (by SDS PAGE) in *Neisseria meningitidis* strains H44/76, M97 250687, BZ10, BZ198, EG328, NGP165, and ATCC 13090.

REMARKS

Claims

Claims 1-18 have been canceled without prejudice or disclaimer of the subject matter therein. Applicant reserves the right to prosecute, in one or more patent applications, the canceled claims, the claims as originally filed, and any other claims supported by the specification.

New claims 19-42 have been introduced. No new matter is added.

Support

Support for the new claims is either obvious, or is as described in the text below. Support for compositions of the isolated polypeptide which include an adjuvant recited in the claims may be found, for example, at page 43, lines 28-29. Support for the hybridization conditions may be found, for example, at page 14, lines 2-7. Support for the recitation of sequence relatedness such as in claim 32 may be found in the specification, for example, at page 47, line 9 through page 48, line 4. Support for the recitation of immune response recognition of a polypeptide of 50 kDa by various *Neisseria meningitidis* strains can be found, for example, at page 60, lines 2-3 and Figures 4 and 5.

International Application No.: PCT/EP00/00425
Attorney Docket No.: BM45353

Closing Remarks

Allowance of the pending claims is respectfully requested.

Respectfully submitted,



Eric A. Meade
Registration No. 42,876
for
Allen Bloom
Registration No. 29,135
Attorney for Applicant

DECHERT
Princeton Pike Corporate Center
PO Box 5218
Princeton, New Jersey 08543-5218
Allen Bloom (609) 620-3214
Eric A. Meade (609) 620-3248
Fax: (609) 620-3259
Attn: Allen Bloom, Esq.
(609) 620-3214

5/10/15

WO 00/43517

PCT/EP00/00425

BASB055 POLYNUCLEOTIDE AND POLYPEPTIDE FROM NEISSERIA MENINGITIDS. USES THEREOF

FIELD OF THE INVENTION

5 This invention relates to polynucleotides, (herein referred to as "BASB055 polynucleotide(s)"), polypeptides encoded by them (referred to herein as "BASB055" or "BASB055 polypeptide(s)"), recombinant materials and methods for their production. In another aspect, the invention relates to methods for using such polypeptides and polynucleotides, including vaccines against bacterial infections. In a further aspect, the
10 invention relates to diagnostic assays for detecting infection of certain pathogens.

BACKGROUND OF THE INVENTION

Neisseria meningitidis (meningococcus) is a Gram-negative bacterium frequently isolated from the human upper respiratory tract. It occasionally causes invasive bacterial diseases
15 such as bacteremia and meningitis. The incidence of meningococcal disease shows geographical seasonal and annual differences (Schwartz, B., Moore, P.S., Broome, C.V.; Clin. Microbiol. Rev. 2 (Supplement), S18-S24, 1989). Most disease in temperate countries is due to strains of serogroup B and varies in incidence from 1-10/100,000/year total population sometimes reaching higher values (Kaczmarek, E.B. (1997), Commun. Dis.
20 Rep. Rev. 7: R55-9, 1995; Scholten, R.J.P.M., Bijlmer, H.A., Poolman, J.T. et al. Clin. Infect. Dis. 16: 237-246, 1993; Cruz, C., Pavez, G., Aguilar, E., et al. Epidemiol. Infect. 105: 119-126, 1990).

Epidemics dominated by serogroup A meningococci, mostly in central Africa, are
25 encountered, sometimes reaching levels up to 1000/100,000/year (Schwartz, B., Moore, P.S., Broome, C.V. Clin. Microbiol. Rev. 2 (Supplement), S18-S24, 1989). Nearly all cases as a whole of meningococcal disease are caused by serogroup A, B, C, W-135 and Y meningococci and a tetravalent A, C, W-135, Y polysaccharide vaccine is available (Armand, J., Arminjon, F., Mynard, M.C., Lafaix, C., J. Biol. Stand. 10: 335-339, 1982).

The polysaccharide vaccines are currently being improved by way of chemical conjugating them to carrier proteins (Lieberman, J.M., Chiu, S.S., Wong, V.K., et al. JAMA 275 : 1499-1503, 1996).

5

A serogroup B vaccine is not available, since the B capsular polysaccharide was found to be nonimmunogenic, most likely because it shares structural similarity to host components (Wyle, F.A., Artenstein, M.S., Brandt, M.L. et al. J. Infect. Dis. 126: 514-522, 1972; Finne, J.M., Leinonen, M., Mäkelä, P.M. Lancet ii.: 355-357, 1983).

10

For many years efforts have been initiated and carried out to develop meningococcal outer membrane based vaccines (de Moraes, J.C., Perkins, B., Camargo, M.C. et al. Lancet 340: 1074-1078, 1992; Bjune, G., Hoiby, E.A. Gronnesby, J.K. et al. 338: 1093-1096, 1991). Such vaccines have demonstrated efficacies from 57% - 85% in older children (>4 years) and adolescents.

15

Many bacterial outer membrane components are present in these vaccines, such as PorA, PorB, Rmp, Opc, Opa, FrpB and the contribution of these components to the observed protection still needs further definition. Other bacterial outer membrane components have been defined by using animal or human antibodies to be potentially relevant to the induction of protective immunity, such as TbpB and NspA (Martin, D., Cadieux, N., Hamel, J., Brodeux, B.R., J. Exp. Med. 185: 1173-1183, 1997; Lissolo, L., Maître-Wilmotte, C., Dumas, p. et al., Inf. Immun. 63: 884-890, 1995). The mechanisms of protective immunity will involve antibody mediated bactericidal activity and opsonophagocytosis.

20

25

A bacteremia animal model has been used to combine all antibody mediated mechanisms (Saukkonen, K., Leinonen, M., Abdillahi, H. Poolman, J. T. Vaccine 7: 325-328, 1989). It is generally accepted that the late complement component mediated bactericidal mechanism is

WO 00/43517

PCT/EP00/00425

crucial for immunity against meningococcal disease (Ross, S.C., Rosenthal P.J., Berberic, H.M., Densen, P. J. Infect. Dis. 155: 1266-1275, 1987).

5 The frequency of *Neisseria meningitidis* infections has risen dramatically in the past few decades. This has been attributed to the emergence of multiply antibiotic resistant strains and an increasing population of people with weakened immune systems. It is no longer uncommon to isolate *Neisseria meningitidis* strains that are resistant to some or all of the standard antibiotics. This phenomenon has created an unmet medical need and demand for new anti-microbial agents, vaccines, drug screening methods, and diagnostic tests for this
10 organism.

SUMMARY OF THE INVENTION

The present invention relates to BASB055, in particular BASB055 polypeptides and
15 BASB055 polynucleotides, recombinant materials and methods for their production. In another aspect, the invention relates to methods for using such polypeptides and polynucleotides, including prevention and treatment of microbial diseases, amongst others. In a further aspect, the invention relates to diagnostic assays for detecting diseases associated with microbial infections and conditions associated with such infections, such
20 as assays for detecting expression or activity of BASB055 polynucleotides or polypeptides.

Various changes and modifications within the spirit and scope of the disclosed invention will become readily apparent to those skilled in the art from reading the following
25 descriptions and from reading the other parts of the present disclosure.

DESCRIPTION OF THE INVENTION

WO 00/43517

PCT/EP00/00425

The BASB055 polypeptide provided in SEQ ID NO:2 is the BASB055 polypeptide from *Neisseria meningitidis* strains ATCC13090.

The invention also provides an immunogenic fragment of a BASB055 polypeptide, that is, a contiguous portion of the BASB055 polypeptide which has the same or substantially the same immunogenic activity as the polypeptide comprising the amino acid sequence of SEQ ID NO:2. That is to say, the fragment (if necessary when coupled to a carrier) is capable of raising an immune response which recognises the BASB055 polypeptide. Such an immunogenic fragment may include, for example, the BASB055 polypeptide lacking an N-terminal leader sequence, and/or a transmembrane domain and/or a C-terminal anchor domain. In a preferred aspect the immunogenic fragment of BASB055 according to the invention comprises substantially all of the extracellular domain of a polypeptide which has at least 85% identity, more preferably at least 90% identity, yet more preferably at least 95% identity, most preferably at least 97-99% identity, to that of SEQ ID NO:2 over the entire length of SEQ ID NO:2.

A fragment is a polypeptide having an amino acid sequence that is entirely the same as part but not all of any amino acid sequence of any polypeptide of the invention. As with BASB055 polypeptides, fragments may be "free-standing," or comprised within a larger polypeptide of which they form a part or region, most preferably as a single continuous region in a single larger polypeptide.

Preferred fragments include, for example, truncation polypeptides having a portion of an amino acid sequence of SEQ ID NO:2 or of variants thereof, such as a continuous series of residues that includes an amino- and/or carboxyl-terminal amino acid sequence. Degradation forms of the polypeptides of the invention produced by or in a host cell, are also preferred. Further preferred are fragments characterized by structural or functional attributes such as fragments that comprise alpha-helix and alpha-helix forming regions, beta-sheet and beta-sheet-forming regions, turn and turn-forming regions, coil and coil-

forming regions, hydrophilic regions, hydrophobic regions, alpha amphipathic regions, beta amphipathic regions, flexible regions, surface-forming regions, substrate binding region, and high antigenic index regions.

- 5 Further preferred fragments include an isolated polypeptide comprising an amino acid sequence having at least 15, 20, 30, 40, 50 or 100 contiguous amino acids from the amino acid sequence of SEQ ID NO:2, or an isolated polypeptide comprising an amino acid sequence having at least 15, 20, 30, 40, 50 or 100 contiguous amino acids truncated or deleted from the amino acid sequence of SEQ ID NO:2.

10

Fragments of the polypeptides of the invention may be employed for producing the corresponding full-length polypeptide by peptide synthesis; therefore, these fragments may be employed as intermediates for producing the full-length polypeptides of the invention.

15

Particularly preferred are variants in which several, 5-10, 1-5, 1-3, 1-2 or 1 amino acids are substituted, deleted, or added in any combination.

20

The polypeptides, or immunogenic fragments, of the invention may be in the form of the "mature" protein or may be a part of a larger protein such as a precursor or a fusion protein. It is often advantageous to include an additional amino acid sequence which contains secretory or leader sequences, pro-sequences, sequences which aid in purification such as multiple histidine residues, or an additional sequence for stability during recombinant production. Furthermore, addition of exogenous polypeptide or

25 lipid tail or polynucleotide sequences to increase the immunogenic potential of the final molecule is also considered.

In one aspect, the invention relates to genetically engineered soluble fusion proteins comprising a polypeptide of the present invention, or a fragment thereof, and various

portions of the constant regions of heavy or light chains of immunoglobulins of various subclasses (IgG, IgM, IgA, IgE). Preferred as an immunoglobulin is the constant part of the heavy chain of human IgG, particularly IgG1, where fusion takes place at the hinge region. In a particular embodiment, the Fc part can be removed simply by incorporation
5 of a cleavage sequence which can be cleaved with blood clotting factor Xa.

Furthermore, this invention relates to processes for the preparation of these fusion proteins by genetic engineering, and to the use thereof for drug screening, diagnosis and therapy. A further aspect of the invention also relates to polynucleotides encoding such
10 fusion proteins. Examples of fusion protein technology can be found in International Patent Application Nos. WO94/29458 and WO94/22914.

The proteins may be chemically conjugated, or expressed as recombinant fusion proteins allowing increased levels to be produced in an expression system as compared
15 to non-fused protein. The fusion partner may assist in providing T helper epitopes (immunological fusion partner), preferably T helper epitopes recognised by humans, or assist in expressing the protein (expression enhancer) at higher yields than the native recombinant protein. Preferably the fusion partner will be both an immunological fusion partner and expression enhancing partner.

20 Fusion partners include protein D from *Haemophilus influenzae* and the non-structural protein from influenzae virus, NS1 (hemagglutinin). Another fusion partner is the protein known as LytA. Preferably the C terminal portion of the molecule is used. LytA is derived from *Streptococcus pneumoniae* which synthesize an N-acetyl-L-
25 alanine amidase, amidase LytA, (coded by the lytA gene {Gene, 43 (1986) page 265-272}) an autolysin that specifically degrades certain bonds in the peptidoglycan backbone. The C-terminal domain of the LytA protein is responsible for the affinity to the choline or to some choline analogues such as DEAE. This property has been exploited for the development of *E.coli* C-LytA expressing plasmids useful for

expression of fusion proteins. Purification of hybrid proteins containing the C-LytA fragment at its amino terminus has been described {Biotechnology: 10, (1992) page 795-798}. It is possible to use the repeat portion of the LytA molecule found in the C terminal end starting at residue 178, for example residues 188 - 305.

5

The present invention also includes variants of the aforementioned polypeptides, that is polypeptides that vary from the referents by conservative amino acid substitutions, whereby a residue is substituted by another with like characteristics. Typical such substitutions are among Ala, Val, Leu and Ile; among Ser and Thr; among the acidic residues Asp and Glu; among Asn and Gln; and among the basic residues Lys and Arg; or aromatic residues Phe and Tyr.

10

Polypeptides of the present invention can be prepared in any suitable manner. Such polypeptides include isolated naturally occurring polypeptides, recombinantly produced polypeptides, synthetically produced polypeptides, or polypeptides produced by a combination of these methods. Means for preparing such polypeptides are well understood in the art.

15

It is most preferred that a polypeptide of the invention is derived from *Neisseria meningitidis*, however, it may preferably be obtained from other organisms of the same taxonomic genus. A polypeptide of the invention may also be obtained, for example, from organisms of the same taxonomic family or order.

20

Polynucleotides

It is an object of the invention to provide polynucleotide that encode BASB055 polypeptide, particularly polynucleotides that encode the polypeptide herein designated BASB055.

25

In a particularly preferred embodiment of the invention the polynucleotide comprises a region encoding BASB055 polypeptide comprising a sequence set out in SEQ ID NO:1 which includes a full length gene, or a variant thereof.

- 5 The BASB055 polynucleotide provided in SEQ ID NO:1 is the BASB055 polynucleotide from *Neisseria meningitidis* strains ATCC13090.

- As a further aspect of the invention there are provided isolated nucleic acid molecules encoding and/or expressing BASB055 polypeptides and polynucleotides, particularly
- 10 *Neisseria meningitidis* BASB055 polypeptides and polynucleotides, including, for example, unprocessed RNAs, ribozyme RNAs, mRNAs, cDNAs, genomic DNAs, B- and Z-DNAs. Further embodiments of the invention include biologically, diagnostically, prophylactically, clinically or therapeutically useful polynucleotides and polypeptides, and variants thereof, and compositions comprising the same.

15

Another aspect of the invention relates to isolated polynucleotides, including at least one full length gene, that encodes a BASB055 polypeptide having a deduced amino acid sequence of SEQ ID NO:2 and polynucleotides closely related thereto and variants thereof.

- 20 In another particularly preferred embodiment of the invention there is a BASB055 polypeptide from *Neisseria meningitidis* comprising or consisting of an amino acid sequence of SEQ ID NO:2 or a variant thereof.

- Using the information provided herein, such as a polynucleotide sequence set out in SEQ ID
- 25 NO:1 a polynucleotide of the invention encoding BASB055 polypeptide may be obtained using standard cloning and screening methods, such as those for cloning and sequencing chromosomal DNA fragments from bacteria using *Neisseria meningitidis* cells as starting material, followed by obtaining a full length clone. For example, to obtain a polynucleotide sequence of the invention, such as a polynucleotide sequence given in SEQ ID NO:1

typically a library of clones of chromosomal DNA of *Neisseria meningitidis* in *E.coli* or some other suitable host is probed with a radiolabeled oligonucleotide, preferably a 17-mer or longer, derived from a partial sequence. Clones carrying DNA identical to that of the probe can then be distinguished using stringent hybridization conditions. By

5 sequencing the individual clones thus identified by hybridization with sequencing primers designed from the original polypeptide or polynucleotide sequence it is then possible to extend the polynucleotide sequence in both directions to determine a full length gene sequence. Conveniently, such sequencing is performed, for example, using denatured double stranded DNA prepared from a plasmid clone. Suitable techniques are described
10 by Maniatis, T., Fritsch, E.F. and Sambrook et al., *MOLECULAR CLONING, A LABORATORY MANUAL*, 2nd Ed.; Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York (1989). (see in particular Screening By Hybridization 1.90 and Sequencing Denatured Double-Stranded DNA Templates 13.70). Direct genomic DNA
15 sequencing may also be performed to obtain a full length gene sequence. Illustrative of the invention, each polynucleotide set out in SEQ ID NO:1 was discovered in a DNA library derived from *Neisseria meningitidis*.

Moreover, each DNA sequence set out in SEQ ID NO:1 contains an open reading frame encoding a protein having about the number of amino acid residues set forth in SEQ ID
20 NO:2 with a deduced molecular weight that can be calculated using amino acid residue molecular weight values well known to those skilled in the art.

The polynucleotide of SEQ ID NO:1, between the start codon at nucleotide number 1 and the stop codon which begins at nucleotide number 1237 of SEQ ID NO:1, encodes the
25 polypeptide of SEQ ID NO:2.

In a further aspect, the present invention provides for an isolated polynucleotide comprising or consisting of:

5 (b) a polynucleotide sequence encoding a polypeptide which has at least 85% identity, more preferably at least 90% identity, yet more preferably at least 95% identity, even more preferably at least 97-99% or 100% exact, to the amino acid sequence of SEQ ID NO:2 over the entire length of SEQ ID NO:2.

15 the sequence of SEQ ID NO:1 or a fragment thereof; and isolating a full-length gene and/or genomic clones containing said polynucleotide sequence.

sequence for a mature polypeptide or a fragment thereof, by itself as well as a coding sequence for a mature polypeptide or a fragment in reading frame with another coding sequence, such as a sequence encoding a leader or secretory sequence, a pre-, or pro- or prepro-protein sequence. The polynucleotide of the invention may also contain at least one non-coding sequence, including for example, but not limited to at least one non-coding 5' and 3' sequence, such as the transcribed but non-translated sequences, termination signals (such as rho-dependent and rho-independent termination signals), ribosome binding sites, Kozak sequences, sequences that stabilize mRNA, introns, and polyadenylation signals. The polynucleotide sequence may also comprise additional coding sequence encoding additional amino acids. For example, a marker sequence that facilitates purification of the

fused polypeptide can be encoded. In certain embodiments of the invention, the marker sequence is a hexa-histidine peptide, as provided in the pQE vector (Qiagen, Inc.) and described in Gentz *et al.*, *Proc. Natl. Acad. Sci., USA* 86: 821-824 (1989), or an HA peptide tag (Wilson *et al.*, *Cell* 37: 767 (1984), both of which may be useful in purifying

5 polypeptide sequence fused to them. Polynucleotides of the invention also include, but are not limited to, polynucleotides comprising a structural gene and its naturally associated sequences that control gene expression.

The nucleotide sequence encoding BASB055 polypeptide of SEQ ID NO:2 may be

10 identical to the polypeptide encoding sequence contained in nucleotides 1 to 1236 of SEQ ID NO:1. Alternatively it may be a sequence, which as a result of the redundancy (degeneracy) of the genetic code, also encodes the polypeptide of SEQ ID NO:2.

The term "polynucleotide encoding a polypeptide" as used herein encompasses

15 polynucleotides that include a sequence encoding a polypeptide of the invention, particularly a bacterial polypeptide and more particularly a polypeptide of the *Neisseria meningitidis* BASB055 having an amino acid sequence set out in SEQ ID NO:2. The term also encompasses polynucleotides that include a single continuous region or discontinuous regions encoding the polypeptide (for example, polynucleotides interrupted by integrated

20 phage, an integrated insertion sequence, an integrated vector sequence, an integrated transposon sequence, or due to RNA editing or genomic DNA reorganization) together with additional regions, that also may contain coding and/or non-coding sequences.

The invention further relates to variants of the polynucleotides described herein that encode

25 variants of a polypeptide having a deduced amino acid sequence of SEQ ID NO:2.

Fragments of polynucleotides of the invention may be used, for example, to synthesize full-length polynucleotides of the invention.

Further particularly preferred embodiments are polynucleotides encoding BASB055 variants, that have the amino acid sequence of BASB055 polypeptide of SEQ ID NO:2 in which several, a few, 5 to 10, 1 to 5, 1 to 3, 2, 1 or no amino acid residues are substituted, modified, deleted and/or added, in any combination. Especially preferred among these are
5 silent substitutions, additions and deletions, that do not alter the properties and activities of BASB055 polypeptide.

Further preferred embodiments of the invention are polynucleotides that are at least 85% identical over their entire length to a polynucleotide encoding BASB055 polypeptide having
10 an amino acid sequence set out in SEQ ID NO:2, and polynucleotides that are complementary to such polynucleotides. In this regard, polynucleotides at least 90% identical over their entire length to the same are particularly preferred, and among these particularly preferred polynucleotides, those with at least 95% are especially preferred. Furthermore, those with at least 97% are highly preferred among those with at least 95%,
15 and among these those with at least 98% and at least 99% are particularly highly preferred, with at least 99% being the more preferred.

Preferred embodiments are polynucleotides encoding polypeptides that retain substantially the same biological function or activity as the mature polypeptide encoded by a DNA of
20 SEQ ID NO:1.

In accordance with certain preferred embodiments of this invention there are provided polynucleotides that hybridize, particularly under stringent conditions, to BASB055 polynucleotide sequences, such as those polynucleotides in SEQ ID NO:1.
25

The invention further relates to polynucleotides that hybridize to the polynucleotide sequences provided herein. In this regard, the invention especially relates to polynucleotides that hybridize under stringent conditions to the polynucleotides described herein. As herein used, the terms "stringent conditions" and "stringent hybridization

conditions" mean hybridization occurring only if there is at least 95% and preferably at least 97% identity between the sequences. A specific example of stringent hybridization conditions is overnight incubation at 42°C in a solution comprising: 50% formamide, 5x SSC (150mM NaCl, 15mM trisodium citrate), 50 mM sodium phosphate (pH7.6), 5x Denhardt's solution, 10% dextran sulfate, and 20 micrograms/ml of denatured, sheared salmon sperm DNA, followed by washing the hybridization support in 0.1x SSC at about 65°C. Hybridization and wash conditions are well known and exemplified in Sambrook, *et al.*, Molecular Cloning: A Laboratory Manual, Second Edition, Cold Spring Harbor, N.Y., (1989), particularly Chapter 11 therein. Solution hybridization may also be used with the polynucleotide sequences provided by the invention.

The invention also provides a polynucleotide consisting of or comprising a polynucleotide sequence obtained by screening an appropriate library containing the complete gene for a polynucleotide sequence set forth in SEQ ID NO:1 under stringent hybridization conditions with a probe having the sequence of said polynucleotide sequence set forth in SEQ ID NO:1 or a fragment thereof; and isolating said polynucleotide sequence. Fragments useful for obtaining such a polynucleotide include, for example, probes and primers fully described elsewhere herein.

As discussed elsewhere herein regarding polynucleotide assays of the invention, for instance, the polynucleotides of the invention, may be used as a hybridization probe for RNA, cDNA and genomic DNA to isolate full-length cDNAs and genomic clones encoding BASB055 and to isolate cDNA and genomic clones of other genes that have a high identity, particularly high sequence identity, to the BASB055 gene. Such probes generally will comprise at least 15 nucleotide residues or base pairs. Preferably, such probes will have at least 30 nucleotide residues or base pairs and may have at least 50 nucleotide residues or base pairs. Particularly preferred probes will have at least 20 nucleotide residues or base pairs and will have less than 30 nucleotide residues or base pairs.

translation systems can also be employed to produce such proteins using RNAs derived from the DNA constructs of the invention.

Recombinant polypeptides of the present invention may be prepared by processes well known in those skilled in the art from genetically engineered host cells comprising expression systems. Accordingly, in a further aspect, the present invention relates to expression systems that comprise a polynucleotide or polynucleotides of the present invention, to host cells which are genetically engineered with such expression systems, and to the production of polypeptides of the invention by recombinant techniques.

For recombinant production of the polypeptides of the invention, host cells can be genetically engineered to incorporate expression systems or portions thereof or polynucleotides of the invention. Introduction of a polynucleotide into the host cell can be effected by methods described in many standard laboratory manuals, such as Davis, *et al.*, *BASIC METHODS IN MOLECULAR BIOLOGY*, (1986) and Sambrook, *et al.*, *MOLECULAR CLONING: A LABORATORY MANUAL*, 2nd Ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. (1989), such as, calcium phosphate transfection, DEAE-dextran mediated transfection, transvection, microinjection, cationic lipid-mediated transfection, electroporation, transduction, scrape loading, ballistic introduction and infection.

Representative examples of appropriate hosts include bacterial cells, such as cells of streptococci, staphylococci, enterococci, *E. coli*, streptomyces, cyanobacteria, *Bacillus subtilis*, *Moraxella catarrhalis*, *Haemophilus influenzae* and *Neisseria meningitidis*; fungal cells, such as cells of a yeast, *Kluveromyces*, *Saccharomyces*, a basidiomycete, *Candida albicans* and *Aspergillus*; insect cells such as cells of *Drosophila* S2 and *Spodoptera* Sf9; animal cells such as CHO, COS, HeLa, C127, 3T3, BHK, 293, CV-1 and Bowes melanoma cells; and plant cells, such as cells of a gymnosperm or angiosperm.

A great variety of expression systems can be used to produce the polypeptides of the invention. Such vectors include, among others, chromosomal-, episomal- and virus-derived vectors, for example, vectors derived from bacterial plasmids, from bacteriophage, from transposons, from yeast episomes, from insertion elements, from yeast chromosomal elements, from viruses such as baculoviruses, papova viruses, such as SV40, vaccinia viruses, adenoviruses, fowl pox viruses, pseudorabies viruses, picornaviruses, retroviruses, and alphaviruses and vectors derived from combinations thereof, such as those derived from plasmid and bacteriophage genetic elements, such as cosmids and phagemids. The expression system constructs may contain control regions that regulate as well as engender expression. Generally, any system or vector suitable to maintain, propagate or express polynucleotides and/or to express a polypeptide in a host may be used for expression in this regard. The appropriate DNA sequence may be inserted into the expression system by any of a variety of well-known and routine techniques, such as, for example, those set forth in Sambrook *et al.*, *MOLECULAR CLONING, A LABORATORY MANUAL*, (*supra*).

In recombinant expression systems in eukaryotes, for secretion of a translated protein into the lumen of the endoplasmic reticulum, into the periplasmic space or into the extracellular environment, appropriate secretion signals may be incorporated into the expressed polypeptide. These signals may be endogenous to the polypeptide or they may be heterologous signals.

Polypeptides of the present invention can be recovered and purified from recombinant cell cultures by well-known methods including ammonium sulfate or ethanol precipitation, acid extraction, anion or cation exchange chromatography, phosphocellulose chromatography, hydrophobic interaction chromatography, affinity chromatography, hydroxylapatite chromatography and lectin chromatography. Most preferably, ion metal affinity chromatography (IMAC) is employed for purification. Well known techniques for refolding proteins may be employed to regenerate active conformation when the polypeptide is denatured during intracellular synthesis, isolation and or purification.

The expression system may also be a recombinant live microorganism, such as a virus or bacterium. The gene of interest can be inserted into the genome of a live recombinant virus or bacterium. Inoculation and *in vivo* infection with this live vector will lead to *in vivo* expression of the antigen and induction of immune responses. Viruses and bacteria used for this purpose are for instance: poxviruses (e.g; vaccinia, fowlpox, canarypox), alphaviruses (Sindbis virus, Semliki Forest Virus, Venezuelan Equine Encephalitis Virus), adenoviruses, adeno-associated virus, picornaviruses (poliovirus, rhinovirus), herpesviruses (varicella zoster virus, etc), Listeria, Salmonella, Shigella, Neisseria, BCG. These viruses and bacteria can be virulent, or attenuated in various ways in order to obtain live vaccines. Such live vaccines also form part of the invention.

Diagnostic, Prognostic, Serotyping and Mutation Assays

This invention is also related to the use of BASB055 polynucleotides and polypeptides of the invention for use as diagnostic reagents. Detection of BASB055 polynucleotides and/or polypeptides in a eukaryote, particularly a mammal, and especially a human, will provide a diagnostic method for diagnosis of disease, staging of disease or response of an infectious organism to drugs. Eukaryotes, particularly mammals, and especially humans, particularly those infected or suspected to be infected with an organism comprising the BASB055 gene or protein, may be detected at the nucleic acid or amino acid level by a variety of well known techniques as well as by methods provided herein.

Polypeptides and polynucleotides for prognosis, diagnosis or other analysis may be obtained from a putatively infected and/or infected individual's bodily materials. Polynucleotides from any of these sources, particularly DNA or RNA, may be used directly for detection or may be amplified enzymatically by using PCR or any other amplification technique prior to analysis. RNA, particularly mRNA, cDNA and genomic DNA may also be used in the same ways. Using amplification, characterization of the species and strain of infectious or resident organism present in an individual, may be made by an analysis of the genotype of a

selected polynucleotide of the organism. Deletions and insertions can be detected by a change in size of the amplified product in comparison to a genotype of a reference sequence selected from a related organism, preferably a different species of the same genus or a different strain of the same species. Point mutations can be identified by hybridizing

5 amplified DNA to labeled BASB055 polynucleotide sequences. Perfectly or significantly matched sequences can be distinguished from imperfectly or more significantly mismatched duplexes by DNase or RNase digestion, for DNA or RNA respectively, or by detecting differences in melting temperatures or renaturation kinetics. Polynucleotide sequence differences may also be detected by alterations in the electrophoretic mobility of

10 polynucleotide fragments in gels as compared to a reference sequence. This may be carried out with or without denaturing agents. Polynucleotide differences may also be detected by direct DNA or RNA sequencing. See, for example, Myers *et al.*, *Science*, 230: 1242 (1985). Sequence changes at specific locations also may be revealed by nuclease protection assays, such as RNase, V1 and S1 protection assay or a chemical cleavage method. See, for

15 example, Cotton *et al.*, *Proc. Natl. Acad. Sci., USA*, 85: 4397-4401 (1985).

In another embodiment, an array of oligonucleotides probes comprising BASB055 nucleotide sequence or fragments thereof can be constructed to conduct efficient screening of, for example, genetic mutations, serotype, taxonomic classification or identification.

20 Array technology methods are well known and have general applicability and can be used to address a variety of questions in molecular genetics including gene expression, genetic linkage, and genetic variability (see, for example, Chee *et al.*, *Science*, 274: 610 (1996)).

Thus in another aspect, the present invention relates to a diagnostic kit which comprises:

- 25 (a) a polynucleotide of the present invention, preferably the nucleotide sequence of SEQ ID NO:1, or a fragment thereof ;
- (b) a nucleotide sequence complementary to that of (a);
- (c) a polypeptide of the present invention, preferably the polypeptide of SEQ ID NO:2 or a fragment thereof; or

(d) an antibody to a polypeptide of the present invention, preferably to the polypeptide of SEQ ID NO:2.

It will be appreciated that in any such kit, (a), (b), (c) or (d) may comprise a substantial
5 component. Such a kit will be of use in diagnosing a disease or susceptibility to a disease, among others.

This invention also relates to the use of polynucleotides of the present invention as diagnostic reagents. Detection of a mutated form of a polynucleotide of the invention,
10 preferable, SEQ ID NO:1, which is associated with a disease or pathogenicity will provide a diagnostic tool that can add to, or define, a diagnosis of a disease, a prognosis of a course of disease, a determination of a stage of disease, or a susceptibility to a disease, which results from under-expression, over-expression or altered expression of the polynucleotide.

Organisms, particularly infectious organisms, carrying mutations in such polynucleotide
15 may be detected at the polynucleotide level by a variety of techniques, such as those described elsewhere herein.

Cells from an organism carrying mutations or polymorphisms (allelic variations) in a polynucleotide and/or polypeptide of the invention may also be detected at the
20 polynucleotide or polypeptide level by a variety of techniques, to allow for serotyping, for example. For example, RT-PCR can be used to detect mutations in the RNA. It is particularly preferred to use RT-PCR in conjunction with automated detection systems, such as, for example, GeneScan. RNA, cDNA or genomic DNA may also be used for the same purpose, PCR. As an example, PCR primers complementary to a polynucleotide encoding
25 BASB055 polypeptide can be used to identify and analyze mutations.

The invention further provides primers with 1, 2, 3 or 4 nucleotides removed from the 5' and/or the 3' end. These primers may be used for, among other things, amplifying BASB055 DNA and/or RNA isolated from a sample derived from an individual, such as a

bodily material. The primers may be used to amplify a polynucleotide isolated from an infected individual, such that the polynucleotide may then be subject to various techniques for elucidation of the polynucleotide sequence. In this way, mutations in the polynucleotide sequence may be detected and used to diagnose and/or prognose the infection or its stage or course, or to serotype and/or classify the infectious agent.

The invention further provides a process for diagnosing disease, preferably bacterial infections, more preferably infections caused by *Neisseria meningitidis*, comprising determining from a sample derived from an individual, such as a bodily material, an increased level of expression of polynucleotide having a sequence of SEQ ID NO:1. Increased or decreased expression of a BASB055 polynucleotide can be measured using any one of the methods well known in the art for the quantitation of polynucleotides, such as, for example, amplification, PCR, RT-PCR, RNase protection, Northern blotting, spectrometry and other hybridization methods.

In addition, a diagnostic assay in accordance with the invention for detecting over-expression of BASB055 polypeptide compared to normal control tissue samples may be used to detect the presence of an infection, for example. Assay techniques that can be used to determine levels of a BASB055 polypeptide, in a sample derived from a host, such as a bodily material, are well-known to those of skill in the art. Such assay methods include radioimmunoassays, competitive-binding assays, Western Blot analysis, antibody sandwich assays, antibody detection and ELISA assays.

The polynucleotides of the invention may be used as components of polynucleotide arrays, preferably high density arrays or grids. These high density arrays are particularly useful for diagnostic and prognostic purposes. For example, a set of spots each comprising a different gene, and further comprising a polynucleotide or polynucleotides of the invention, may be used for probing, such as using hybridization or nucleic acid amplification, using a probe obtained or derived from a bodily sample,

to determine the presence of a particular polynucleotide sequence or related sequence in an individual. Such a presence may indicate the presence of a pathogen, particularly *Neisseria meningitidis*, and may be useful in diagnosing and/or prognosing disease or a course of disease. A grid comprising a number of variants of the polynucleotide sequence of SEQ ID NO:1 are preferred. Also preferred is a grid comprising a number of variants of a polynucleotide sequence encoding the polypeptide sequence of SEQ ID NO:2.

Antibodies

- 10 The polypeptides and polynucleotides of the invention or variants thereof, or cells expressing the same can be used as immunogens to produce antibodies immunospecific for such polypeptides or polynucleotides respectively.

- In certain preferred embodiments of the invention there are provided antibodies against
15 BASB055 polypeptides or polynucleotides.

- Antibodies generated against the polypeptides or polynucleotides of the invention can be obtained by administering the polypeptides and/or polynucleotides of the invention, or epitope-bearing fragments of either or both, analogues of either or both, or cells expressing
20 either or both, to an animal, preferably a nonhuman, using routine protocols. For preparation of monoclonal antibodies, any technique known in the art that provides antibodies produced by continuous cell line cultures can be used. Examples include various techniques, such as those in Kohler, G. and Milstein, C., *Nature* 256: 495-497 (1975); Kozbor *et al.*, *Immunology Today* 4: 72 (1983); Cole *et al.*, pg. 77-96 in *MONOCLONAL*
25 *ANTIBODIES AND CANCER THERAPY*, Alan R. Liss, Inc. (1985).

Techniques for the production of single chain antibodies (U.S. Patent No. 4,946,778) can be adapted to produce single chain antibodies to polypeptides or polynucleotides of this invention. Also, transgenic mice, or other organisms or animals, such as other mammals,

may be used to express humanized antibodies immunospecific to the polypeptides or polynucleotides of the invention.

Alternatively, phage display technology may be utilized to select antibody genes with
5 binding activities towards a polypeptide of the invention either from repertoires of PCR amplified v-genes of lymphocytes from humans screened for possessing anti-BASB055 or from naive libraries (McCafferty, *et al.*, (1990), *Nature* 348, 552-554; Marks, *et al.*, (1992) *Biotechnology* 10, 779-783). The affinity of these antibodies can also be improved by, for example, chain shuffling (Clackson *et al.*, (1991) *Nature* 352: 628).

10 The above-described antibodies may be employed to isolate or to identify clones expressing the polypeptides or polynucleotides of the invention to purify the polypeptides or polynucleotides by, for example, affinity chromatography.

15 Thus, among others, antibodies against BASB055-polypeptide or BASB055-polynucleotide may be employed to treat infections, particularly bacterial infections.

Polypeptide variants include antigenically, epitopically or immunologically equivalent variants form a particular aspect of this invention.

20 Preferably, the antibody or variant thereof is modified to make it less immunogenic in the individual. For example, if the individual is human the antibody may most preferably be "humanized," where the complementarity determining region or regions of the hybridoma-derived antibody has been transplanted into a human monoclonal antibody, for example
25 as described in Jones *et al.* (1986), *Nature* 321, 522-525 or Tempest *et al.*, (1991) *Biotechnology* 9, 266-273.

Antagonists and Agonists - Assays and Molecules

Polypeptides and polynucleotides of the invention may also be used to assess the binding of small molecule substrates and ligands in, for example, cells, cell-free preparations, chemical libraries, and natural product mixtures. These substrates and ligands may be natural substrates and ligands or may be structural or functional mimetics. See, *e.g.*, Coligan *et al.*,
5 *Current Protocols in Immunology* 1(2): Chapter 5 (1991).

The screening methods may simply measure the binding of a candidate compound to the polypeptide or polynucleotide, or to cells or membranes bearing the polypeptide or polynucleotide, or a fusion protein of the polypeptide by means of a label directly or
10 indirectly associated with the candidate compound. Alternatively, the screening method may involve competition with a labeled competitor. Further, these screening methods may test whether the candidate compound results in a signal generated by activation or inhibition of the polypeptide or polynucleotide, using detection systems appropriate to the cells comprising the polypeptide or polynucleotide. Inhibitors of activation are generally
15 assayed in the presence of a known agonist and the effect on activation by the agonist by the presence of the candidate compound is observed. Constitutively active polypeptide and/or constitutively expressed polypeptides and polynucleotides may be employed in screening methods for inverse agonists or inhibitors, in the absence of an agonist or inhibitor, by testing whether the candidate compound results in inhibition of activation of
20 the polypeptide or polynucleotide, as the case may be. Further, the screening methods may simply comprise the steps of mixing a candidate compound with a solution containing a polypeptide or polynucleotide of the present invention, to form a mixture, measuring BASB055 polypeptide and/or polynucleotide activity in the mixture, and comparing the BASB055 polypeptide and/or polynucleotide activity of the mixture to a
25 standard. Fusion proteins, such as those made from Fc portion and BASB055 polypeptide, as hereinbefore described, can also be used for high-throughput screening assays to identify antagonists of the polypeptide of the present invention, as well as of phylogenetically and and/or functionally related polypeptides (see D. Bennett *et al.*, J Mol

Recognition, 8:52-58 (1995); and K. Johanson *et al.*, J Biol Chem, 270(16):9459-9471 (1995)).

The polynucleotides, polypeptides and antibodies that bind to and/or interact with a
5 polypeptide of the present invention may also be used to configure screening methods for
detecting the effect of added compounds on the production of mRNA and/or polypeptide
in cells. For example, an ELISA assay may be constructed for measuring secreted or cell
associated levels of polypeptide using monoclonal and polyclonal antibodies by standard
methods known in the art. This can be used to discover agents which may inhibit or
10 enhance the production of polypeptide (also called antagonist or agonist, respectively)
from suitably manipulated cells or tissues.

The invention also provides a method of screening compounds to identify those which
enhance (agonist) or block (antagonist) the action of BASB055 polypeptides or
15 polynucleotides, particularly those compounds that are bacteristatic and/or bactericidal. The
method of screening may involve high-throughput techniques. For example, to screen for
agonists or antagonists, a synthetic reaction mix, a cellular compartment, such as a
membrane, cell envelope or cell wall, or a preparation of any thereof, comprising BASB055
polypeptide and a labeled substrate or ligand of such polypeptide is incubated in the absence
20 or the presence of a candidate molecule that may be a BASB055 agonist or antagonist. The
ability of the candidate molecule to agonize or antagonize the BASB055 polypeptide is
reflected in decreased binding of the labeled ligand or decreased production of product from
such substrate. Molecules that bind gratuitously, *i.e.*, without inducing the effects of
BASB055 polypeptide are most likely to be good antagonists. Molecules that bind well
25 and, as the case may be, increase the rate of product production from substrate, increase
signal transduction, or increase chemical channel activity are agonists. Detection of the rate
or level of, as the case may be, production of product from substrate, signal transduction, or
chemical channel activity may be enhanced by using a reporter system. Reporter systems
that may be useful in this regard include but are not limited to colorimetric, labeled substrate

converted into product, a reporter gene that is responsive to changes in BASB055 polynucleotide or polypeptide activity, and binding assays known in the art.

Another example of an assay for BASB055 agonists is a competitive assay that combines
5 BASB055 and a potential agonist with BASB055-binding molecules, recombinant BASB055 binding molecules, natural substrates or ligands, or substrate or ligand mimetics, under appropriate conditions for a competitive inhibition assay. BASB055 can be labeled, such as by radioactivity or a colorimetric compound, such that the number of BASB055 molecules bound to a binding molecule or converted to product can be determined
10 accurately to assess the effectiveness of the potential antagonist.

Potential antagonists include, among others, small organic molecules, peptides, polypeptides and antibodies that bind to a polynucleotide and/or polypeptide of the invention and thereby inhibit or extinguish its activity or expression. Potential antagonists
15 also may be small organic molecules, a peptide, a polypeptide such as a closely related protein or antibody that binds the same sites on a binding molecule, such as a binding molecule, without inducing BASB055-induced activities, thereby preventing the action or expression of BASB055 polypeptides and/or polynucleotides by excluding BASB055 polypeptides and/or polynucleotides from binding.

20

Potential antagonists include a small molecule that binds to and occupies the binding site of the polypeptide thereby preventing binding to cellular binding molecules, such that normal biological activity is prevented. Examples of small molecules include but are not limited to small organic molecules, peptides or peptide-like molecules. Other potential antagonists
25 include antisense molecules (see Okano, *J. Neurochem.* 56: 560 (1991); *OLIGODEOXYNUCLEOTIDES AS ANTISENSE INHIBITORS OF GENE EXPRESSION*, CRC Press, Boca Raton, FL (1988), for a description of these molecules). Preferred potential antagonists include compounds related to and variants of BASB055.

In a further aspect, the present invention relates to genetically engineered soluble fusion proteins comprising a polypeptide of the present invention, or a fragment thereof, and various portions of the constant regions of heavy or light chains of immunoglobulins of various subclasses (IgG, IgM, IgA, IgE). Preferred as an immunoglobulin is the constant part of the heavy chain of human IgG, particularly IgG1, where fusion takes place at the hinge region. In a particular embodiment, the Fc part can be removed simply by incorporation of a cleavage sequence which can be cleaved with blood clotting factor Xa. Furthermore, this invention relates to processes for the preparation of these fusion proteins by genetic engineering, and to the use thereof for drug screening, diagnosis and therapy. A further aspect of the invention also relates to polynucleotides encoding such fusion proteins. Examples of fusion protein technology can be found in International Patent Application Nos. WO94/29458 and WO94/22914.

Each of the polynucleotide sequences provided herein may be used in the discovery and development of antibacterial compounds. The encoded protein, upon expression, can be used as a target for the screening of antibacterial drugs. Additionally, the polynucleotide sequences encoding the amino terminal regions of the encoded protein or Shine-Delgarno or other translation facilitating sequences of the respective mRNA can be used to construct antisense sequences to control the expression of the coding sequence of interest.

The invention also provides the use of the polypeptide, polynucleotide, agonist or antagonist of the invention to interfere with the initial physical interaction between a pathogen or pathogens and a eukaryotic, preferably mammalian, host responsible for sequelae of infection. In particular, the molecules of the invention may be used: in the prevention of adhesion of bacteria, in particular gram positive and/or gram negative bacteria, to eukaryotic, preferably mammalian, extracellular matrix proteins on in-dwelling devices or to extracellular matrix proteins in wounds; to block bacterial adhesion between eukaryotic, preferably mammalian, extracellular matrix proteins and bacterial BASB055 proteins that mediate tissue damage and/or; to block the normal progression of

Alternatively, peptide mimotopes may be identified using antibodies which are capable themselves of binding to the polypeptides of the present invention using techniques such as phage display technology (EP 0 552 267 B1). This technique, generates a large number of peptide sequences which mimic the structure of the native peptides and are, therefore, capable of binding to anti-native peptide antibodies, but may not necessarily themselves share significant sequence homology to the native polypeptide.

Vaccines

Another aspect of the invention relates to a method for inducing an immunological response in an individual, particularly a mammal, preferably humans, which comprises inoculating the individual with BASB055 polynucleotide and/or polypeptide, or a fragment or variant thereof, adequate to produce antibody and/ or T cell immune response to protect said individual from infection, particularly bacterial infection and most particularly *Neisseria meningitidis* infection. Also provided are methods whereby such immunological response slows bacterial replication. Yet another aspect of the invention relates to a method of inducing immunological response in an individual which comprises delivering to such individual a nucleic acid vector, sequence or ribozyme to direct expression of BASB055 polynucleotide and/or polypeptide, or a fragment or a variant thereof, for expressing BASB055 polynucleotide and/or polypeptide, or a fragment or a variant thereof *in vivo* in order to induce an immunological response, such as, to produce antibody and/ or T cell immune response, including, for example, cytokine-producing T cells or cytotoxic T cells, to protect said individual, preferably a human, from disease, whether that disease is already established within the individual or not. One example of administering the gene is by accelerating it into the desired cells as a coating on particles or otherwise. Such nucleic acid vector may comprise DNA, RNA, a ribozyme, a modified nucleic acid, a DNA/RNA hybrid, a DNA-protein complex or an RNA-protein complex.

A further aspect of the invention relates to an immunological composition that when introduced into an individual, preferably a human, capable of having induced within it an immunological response, induces an immunological response in such individual to a BASB055 polynucleotide and/or polypeptide encoded therefrom, wherein the

5 composition comprises a recombinant BASB055 polynucleotide and/or polypeptide encoded therefrom and/or comprises DNA and/or RNA which encodes and expresses an antigen of said BASB055 polynucleotide, polypeptide encoded therefrom, or other polypeptide of the invention. The immunological response may be used therapeutically or prophylactically and may take the form of antibody immunity and/or cellular

10 immunity, such as cellular immunity arising from CTL or CD4+ T cells.

A BASB055 polypeptide or a fragment thereof may be fused with co-protein or chemical moiety which may or may not by itself produce antibodies, but which is capable of stabilizing the first protein and producing a fused or modified protein which will have

15 antigenic and/or immunogenic properties, and preferably protective properties. Thus fused recombinant protein, preferably further comprises an antigenic co-protein, such as lipoprotein D from *Haemophilus influenzae*, Glutathione-S-transferase (GST) or beta-galactosidase, or any other relatively large co-protein which solubilizes the protein and facilitates production and purification thereof. Moreover, the co-protein may act as an

20 adjuvant in the sense of providing a generalized stimulation of the immune system of the organism receiving the protein. The co-protein may be attached to either the amino- or carboxy-terminus of the first protein.

In a vaccine composition according to the invention, a BASB055 polypeptide and/or

25 polynucleotide, or a fragment, or a mimotope, or a variant thereof may be present in a vector, such as the live recombinant vectors described above for example live bacterial vectors.

with potential secondary structure involved in regulation, as well as other types of regulatory motifs or sequences. This sequence is a further aspect of the invention.

This sequence information allows the modulation of the natural expression of the
5 BASB055 gene. The upregulation of the gene expression may be accomplished by
altering the promoter, the shine-dalgarno sequence, potential repressor or operator
elements, or any other elements involved. Likewise, downregulation of expression can be
achieved by similar types of modification. Alternatively, by changing phase variation
10 sequences, the expression of the gene can be put under phase variation control, or it may
be uncoupled from this regulation. In another approach, the expression of the gene can be
put under the control of one or more inducible elements allowing regulated expression.
Examples of such regulation include, but are not limited to, induction by temperature
shift, addition of inductor substrates like selected carbohydrates or their derivatives, trace
elements, vitamins, co-factors, metal ions, etc.

15
Such modifications as described above can be introduced by several different means. The
modification of sequences involved in gene expression can be carried out *in vivo* by
random mutagenesis followed by selection for the desired phenotype. Another approach
consists in isolating the region of interest and modifying it by random mutagenesis, or
20 site-directed replacement, insertion or deletion mutagenesis. The modified region can then
be reintroduced into the bacterial genome by homologous recombination, and the effect
on gene expression can be assessed. In another approach, the sequence knowledge of the
region of interest can be used to replace or delete all or part of the natural regulatory
sequences. In this case, the regulatory region targeted is isolated and modified so as to
25 contain the regulatory elements from another gene, a combination of regulatory elements
from different genes, a synthetic regulatory region, or any other regulatory region, or to
delete selected parts of the wild-type regulatory sequences. These modified sequences can
then be reintroduced into the bacterium via homologous recombination into the genome.
A non-exhaustive list of preferred promoters that could be used for up-regulation of gene

Thus, the invention provides a BASB055 polypeptide, in a modified bacterial bleb. The invention further provides modified host cells capable of producing the non-live membrane-based bleb vectors. The invention further provides nucleic acid vectors comprising the BASB055 gene having a modified upstream region containing a heterologous regulatory element.

Further provided by the invention are processes to prepare the host cells and bacterial blebs according to the invention.

Also provided by this invention are compositions, particularly vaccine compositions, and methods comprising the polypeptides and/or polynucleotides of the invention and immunostimulatory DNA sequences, such as those described in Sato, Y. *et al.* Science 273: 352 (1996).

Also, provided by this invention are methods using the described polynucleotide or particular fragments thereof, which have been shown to encode non-variable regions of bacterial cell surface proteins, in polynucleotide constructs used in such genetic immunization experiments in animal models of infection with *Neisseria meningitidis*.

Such experiments will be particularly useful for identifying protein epitopes able to provoke a prophylactic or therapeutic immune response. It is believed that this approach will allow for the subsequent preparation of monoclonal antibodies of particular value, derived from the requisite organ of the animal successfully resisting or clearing infection, for the development of prophylactic agents or therapeutic treatments of bacterial infection, particularly *Neisseria meningitidis* infection, in mammals, particularly humans.

The invention also includes a vaccine formulation which comprises an immunogenic recombinant polypeptide and/or polynucleotide of the invention together with a suitable carrier, such as a pharmaceutically acceptable carrier. Since the polypeptides and

polynucleotides may be broken down in the stomach, each is preferably administered parenterally, including, for example, administration that is subcutaneous, intramuscular, intravenous, or intradermal. Formulations suitable for parenteral administration include aqueous and non-aqueous sterile injection solutions which may contain anti-oxidants, buffers, bacteristatic compounds and solutes which render the formulation isotonic with the bodily fluid, preferably the blood, of the individual; and aqueous and non-aqueous sterile suspensions which may include suspending agents or thickening agents. The formulations may be presented in unit-dose or multi-dose containers, for example, sealed ampoules and vials and may be stored in a freeze-dried condition requiring only the addition of the sterile liquid carrier immediately prior to use.

The vaccine formulation of the invention may also include adjuvant systems for enhancing the immunogenicity of the formulation. Preferably the adjuvant system raises preferentially a TH1 type of response.

An immune response may be broadly distinguished into two extreme categories, being a humoral or cell mediated immune responses (traditionally characterised by antibody and cellular effector mechanisms of protection respectively). These categories of response have been termed TH1-type responses (cell-mediated response), and TH2-type immune responses (humoral response).

Extreme TH1-type immune responses may be characterised by the generation of antigen specific, haplotype restricted cytotoxic T lymphocytes, and natural killer cell responses. In mice TH1-type responses are often characterised by the generation of antibodies of the IgG2a subtype, whilst in the human these correspond to IgG1 type antibodies. TH2-type immune responses are characterised by the generation of a broad range of immunoglobulin isotypes including in mice IgG1, IgA, and IgM.

It can be considered that the driving force behind the development of these two types of immune responses are cytokines. High levels of TH1-type cytokines tend to favour the induction of cell mediated immune responses to the given antigen, whilst high levels of TH2-type cytokines tend to favour the induction of humoral immune responses to the antigen.

The distinction of TH1 and TH2-type immune responses is not absolute. In reality an individual will support an immune response which is described as being predominantly TH1 or predominantly TH2. However, it is often convenient to consider the families of cytokines in terms of that described in murine CD4 +ve T cell clones by Mosmann and Coffman (*Mosmann, T.R. and Coffman, R.L. (1989) TH1 and TH2 cells: different patterns of lymphokine secretion lead to different functional properties. Annual Review of Immunology, 7, p145-173*). Traditionally, TH1-type responses are associated with the production of the INF- γ and IL-2 cytokines by T-lymphocytes. Other cytokines often directly associated with the induction of TH1-type immune responses are not produced by T-cells, such as IL-12. In contrast, TH2- type responses are associated with the secretion of IL-4, IL-5, IL-6 and IL-13.

It is known that certain vaccine adjuvants are particularly suited to the stimulation of either TH1 or TH2 - type cytokine responses. Traditionally the best indicators of the TH1:TH2 balance of the immune response after a vaccination or infection includes direct measurement of the production of TH1 or TH2 cytokines by T lymphocytes *in vitro* after restimulation with antigen, and/or the measurement of the IgG1:IgG2a ratio of antigen specific antibody responses.

Thus, a TH1-type adjuvant is one which preferentially stimulates isolated T-cell populations to produce high levels of TH1-type cytokines when re-stimulated with antigen *in vitro*, and promotes development of both CD8+ cytotoxic T lymphocytes and antigen specific immunoglobulin responses associated with TH1-type isotype.

Adjuvants which are capable of preferential stimulation of the TH1 cell response are described in International Patent Application No. WO 94/00153 and WO 95/17209.

5 3 De-O-acylated monophosphoryl lipid A (3D-MPL) is one such adjuvant. This is known from GB 2220211 (Ribi). Chemically it is a mixture of 3 De-O-acylated monophosphoryl lipid A with 4, 5 or 6 acylated chains and is manufactured by Ribi Immunochem, Montana. A preferred form of 3 De-O-acylated monophosphoryl lipid A is disclosed in European Patent 0 689 454 B1 (SmithKline Beecham Biologicals SA).

10

Preferably, the particles of 3D-MPL are small enough to be sterile filtered through a 0.22micron membrane (European Patent number 0 689 454).

3D-MPL will be present in the range of 10µg - 100µg preferably 25-50µg per dose wherein the antigen will typically be present in a range 2-50µg per dose.

15

Another preferred adjuvant comprises QS21, an Hplc purified non-toxic fraction derived from the bark of Quillaja Saponaria Molina. Optionally this may be admixed with 3 De-O-acylated monophosphoryl lipid A (3D-MPL), optionally together with a carrier.

20

The method of production of QS21 is disclosed in US patent No. 5,057,540.

Non-reactogenic adjuvant formulations containing QS21 have been described previously (WO 96/33739). Such formulations comprising QS21 and cholesterol have
25 been shown to be successful TH1 stimulating adjuvants when formulated together with an antigen.

Further adjuvants which are preferential stimulators of TH1 cell response include immunomodulatory oligonucleotides, for example unmethylated CpG sequences as disclosed in WO 96/02555.

- 5 Combinations of different TH1 stimulating adjuvants, such as those mentioned hereinabove, are also contemplated as providing an adjuvant which is a preferential stimulator of TH1 cell response. For example, QS21 can be formulated together with 3D-MPL. The ratio of QS21 : 3D-MPL will typically be in the order of 1 : 10 to 10 : 1; preferably 1:5 to 5 : 1 and often substantially 1 : 1. The preferred range for optimal
10 synergy is 2.5 : 1 to 1 : 1 3D-MPL: QS21.

Preferably a carrier is also present in the vaccine composition according to the invention. The carrier may be an oil in water emulsion, or an aluminium salt, such as aluminium phosphate or aluminium hydroxide.

15

A preferred oil-in-water emulsion comprises a metabolisable oil, such as squalene, alpha tocopherol and Tween 80. In a particularly preferred aspect the antigens in the vaccine composition according to the invention are combined with QS21 and 3D-MPL in such an emulsion. Additionally the oil in water emulsion may contain span 85 and/or
20 lecithin and/or tricaprylin.

Typically for human administration QS21 and 3D-MPL will be present in a vaccine in the range of 1µg - 200µg, such as 10-100µg, preferably 10µg - 50µg per dose.

- Typically the oil in water will comprise from 2 to 10% squalene, from 2 to 10% alpha
25 tocopherol and from 0.3 to 3% tween 80. Preferably the ratio of squalene: alpha tocopherol is equal to or less than 1 as this provides a more stable emulsion. Span 85 may also be present at a level of 1%. In some cases it may be advantageous that the vaccines of the present invention will further contain a stabiliser.

or more of the ingredients of the aforementioned compositions of the invention.

Polypeptides, polynucleotides and other compounds of the present invention may be employed alone or in conjunction with other compounds, such as therapeutic compounds.

- 5 The composition will be adapted to the route of administration, for instance by a systemic or an oral route. Preferred forms of systemic administration include injection, typically by intravenous injection. Other injection routes, such as subcutaneous, intramuscular, or intraperitoneal, can be used. Alternative means for systemic administration include transmucosal and transdermal administration using penetrants such as bile salts or fusidic
- 10 acids or other detergents. In addition, if a polypeptide or other compounds of the present invention can be formulated in an enteric or an encapsulated formulation, oral administration may also be possible. Administration of these compounds may also be topical and/or localized, in the form of salves, pastes, gels, solutions, powders and the like.
- 15 For administration to mammals, and particularly humans, it is expected that the daily dosage level of the active agent will be from 0.01 mg/kg to 10 mg/kg, typically around 1 mg/kg. The physician in any event will determine the actual dosage which will be most suitable for an individual and will vary with the age, weight and response of the particular individual. The above dosages are exemplary of the average case. There can, of course,
- 20 be individual instances where higher or lower dosage ranges are merited, and such are within the scope of this invention.

- The dosage range required depends on the choice of peptide, the route of administration, the nature of the formulation, the nature of the subject's condition, and the judgment of the
- 25 attending practitioner. Suitable dosages, however, are in the range of 0.1-100 µg/kg of subject.

A vaccine composition is conveniently in injectable form. Conventional adjuvants may be employed to enhance the immune response. A suitable unit dose for vaccination is 0.5-5

microgram/kg of antigen, and such dose is preferably administered 1-3 times and with an interval of 1-3 weeks. With the indicated dose range, no adverse toxicological effects will be observed with the compounds of the invention which would preclude their administration to suitable individuals.

5

Wide variations in the needed dosage, however, are to be expected in view of the variety of compounds available and the differing efficiencies of various routes of administration. For example, oral administration would be expected to require higher dosages than administration by intravenous injection. Variations in these dosage levels can be adjusted using standard empirical routines for optimization, as is well understood in the art.

10

Sequence Databases, Sequences in a Tangible Medium, and Algorithms

Polynucleotide and polypeptide sequences form a valuable information resource with which to determine their 2- and 3-dimensional structures as well as to identify further sequences of similar homology. These approaches are most easily facilitated by storing the sequence in a computer readable medium and then using the stored data in a known macromolecular structure program or to search a sequence database using well known searching tools, such as the GCG program package.

15

20

Also provided by the invention are methods for the analysis of character sequences or strings, particularly genetic sequences or encoded protein sequences. Preferred methods of sequence analysis include, for example, methods of sequence homology analysis, such as identity and similarity analysis, DNA, RNA and protein structure analysis, sequence assembly, cladistic analysis, sequence motif analysis, open reading frame determination, nucleic acid base calling, codon usage analysis, nucleic acid base trimming, and sequencing chromatogram peak analysis.

25

A computer based method is provided for performing homology identification. This method comprises the steps of: providing a first polynucleotide sequence comprising the sequence of a polynucleotide of the invention in a computer readable medium; and comparing said first polynucleotide sequence to at least one second polynucleotide or polypeptide sequence to identify homology.

A computer based method is also provided for performing homology identification, said method comprising the steps of: providing a first polypeptide sequence comprising the sequence of a polypeptide of the invention in a computer readable medium; and comparing said first polypeptide sequence to at least one second polynucleotide or polypeptide sequence to identify homology.

All publications and references, including but not limited to patents and patent applications, cited in this specification are herein incorporated by reference in their entirety as if each individual publication or reference were specifically and individually indicated to be incorporated by reference herein as being fully set forth. Any patent application to which this application claims priority is also incorporated by reference herein in its entirety in the manner described above for publications and references.

DEFINITIONS

"Identity," as known in the art, is a relationship between two or more polypeptide sequences or two or more polynucleotide sequences, as the case may be, as determined by comparing the sequences. In the art, "identity" also means the degree of sequence relatedness between polypeptide or polynucleotide sequences, as the case may be, as determined by the match between strings of such sequences. "Identity" can be readily calculated by known methods, including but not limited to those described in (*Computational Molecular Biology*, Lesk, A.M., ed., Oxford University Press, New York, 1988; *Biocomputing: Informatics and Genome Projects*, Smith, D.W., ed., Academic Press, New York, 1993;

Computer Analysis of Sequence Data, Part I, Griffin, A.M., and Griffin, H.G., eds., Humana Press, New Jersey, 1994; *Sequence Analysis in Molecular Biology*, von Heine, G., Academic Press, 1987; and *Sequence Analysis Primer*, Gribskov, M. and Devereux, J., eds., M Stockton Press, New York, 1991; and Carillo, H., and Lipman, D., SIAM J.

- 5 *Applied Math.*, 48: 1073 (1988). Methods to determine identity are designed to give the largest match between the sequences tested. Moreover, methods to determine identity are codified in publicly available computer programs. Computer program methods to determine identity between two sequences include, but are not limited to, the GAP program in the GCG program package (Devereux, J., et al., *Nucleic Acids Research* 12(1):
- 10 387 (1984)), BLASTP, BLASTN (Altschul, S.F. et al., *J. Mol. Biol.* 215: 403-410 (1990), and FASTA(Pearson and Lipman Proc. Natl. Acad. Sci. USA 85; 2444-2448 (1988). The BLAST family of programs is publicly available from NCBI and other sources (*BLAST Manual*, Altschul, S., et al., NCBI NLM NIH Bethesda, MD 20894; Altschul, S., et al., *J. Mol. Biol.* 215: 403-410 (1990). The well known Smith Waterman algorithm
- 15 may also be used to determine identity.

Parameters for polypeptide sequence comparison include the following:

Algorithm: Needleman and Wunsch, *J. Mol Biol.* 48: 443-453 (1970)

Comparison matrix: BLOSSUM62 from Henikoff and Henikoff,

- 20 Proc. Natl. Acad. Sci. USA. 89:10915-10919 (1992)

Gap Penalty: 8

Gap Length Penalty: 2

A program useful with these parameters is publicly available as the "gap" program from Genetics Computer Group, Madison WI. The aforementioned parameters are the default

25 parameters for peptide comparisons (along with no penalty for end gaps).

Parameters for polynucleotide comparison include the following:

Algorithm: Needleman and Wunsch, *J. Mol Biol.* 48: 443-453 (1970)

Comparison matrix: matches = +10, mismatch = 0

WO 00/43517

PCT/EP00/00425

Gap Penalty: 50

Gap Length Penalty: 3

Available as: The "gap" program from Genetics Computer Group, Madison WI. These are the default parameters for nucleic acid comparisons.

5

A preferred meaning for "identity" for polynucleotides and polypeptides, as the case may be, are provided in (1) and (2) below.

- (1) Polynucleotide embodiments further include an isolated polynucleotide
- 10 comprising a polynucleotide sequence having at least a 50, 60, 70, 80, 85, 90, 95, 97 or 100% identity to the reference sequence of SEQ ID NO:1, wherein said polynucleotide sequence may be identical to the reference sequence of SEQ ID NO:1 or may include up to a certain integer number of nucleotide alterations as compared to the reference sequence, wherein said alterations are selected from the group consisting of at least one
- 15 nucleotide deletion, substitution, including transition and transversion, or insertion, and wherein said alterations may occur at the 5' or 3' terminal positions of the reference nucleotide sequence or anywhere between those terminal positions, interspersed either individually among the nucleotides in the reference sequence or in one or more contiguous groups within the reference sequence, and wherein said number of nucleotide
- 20 alterations is determined by multiplying the total number of nucleotides in SEQ ID NO:1 by the integer defining the percent identity divided by 100 and then subtracting that product from said total number of nucleotides in SEQ ID NO:1, or:

$$n_n \leq x_n - (x_n \bullet y),$$

25

wherein n_n is the number of nucleotide alterations, x_n is the total number of nucleotides in SEQ ID NO:1, y is 0.50 for 50%, 0.60 for 60%, 0.70 for 70%, 0.80 for 80%, 0.85 for 85%, 0.90 for 90%, 0.95 for 95%, 0.97 for 97% or 1.00 for 100%, and \bullet is the symbol for the multiplication operator, and wherein any non-integer product of x_n and y is rounded

down to the nearest integer prior to subtracting it from x_n . Alterations of a polynucleotide sequence encoding the polypeptide of SEQ ID NO:2 may create nonsense, missense or frameshift mutations in this coding sequence and thereby alter the polypeptide encoded by the polynucleotide following such alterations.

5

By way of example, a polynucleotide sequence of the present invention may be identical to the reference sequence of SEQ ID NO:1, that is it may be 100% identical, or it may include up to a certain integer number of nucleic acid alterations as compared to the reference sequence such that the percent identity is less than 100% identity. Such

10 alterations are selected from the group consisting of at least one nucleic acid deletion, substitution, including transition and transversion, or insertion, and wherein said alterations may occur at the 5' or 3' terminal positions of the reference polynucleotide sequence or anywhere between those terminal positions, interspersed either individually among the nucleic acids in the reference sequence or in one or more contiguous groups
15 within the reference sequence. The number of nucleic acid alterations for a given percent identity is determined by multiplying the total number of nucleic acids in SEQ ID NO:1 by the integer defining the percent identity divided by 100 and then subtracting that product from said total number of nucleic acids in SEQ ID NO:1, or:

$$20 \quad n_n \leq x_n - (x_n \bullet y),$$

wherein n_n is the number of nucleic acid alterations, x_n is the total number of nucleic acids in SEQ ID NO:1, y is, for instance 0.70 for 70%, 0.80 for 80%, 0.85 for 85% etc., \bullet is the symbol for the multiplication operator, and wherein any non-integer product of x_n
25 and y is rounded down to the nearest integer prior to subtracting it from x_n .

(2) Polypeptide embodiments further include an isolated polypeptide comprising a polypeptide having at least a 50, 60, 70, 80, 85, 90, 95, 97 or 100% identity to a polypeptide reference sequence of SEQ ID NO:2, wherein said polypeptide sequence may

be identical to the reference sequence of SEQ ID NO:2 or may include up to a certain integer number of amino acid alterations as compared to the reference sequence, wherein said alterations are selected from the group consisting of at least one amino acid deletion, substitution, including conservative and non-conservative substitution, or insertion, and
 5 wherein said alterations may occur at the amino- or carboxy-terminal positions of the reference polypeptide sequence or anywhere between those terminal positions, interspersed either individually among the amino acids in the reference sequence or in one or more contiguous groups within the reference sequence, and wherein said number of amino acid alterations is determined by multiplying the total number of amino acids in
 10 SEQ ID NO:2 by the integer defining the percent identity divided by 100 and then subtracting that product from said total number of amino acids in SEQ ID NO:2, or:

$$n_a \leq x_a - (x_a \bullet y),$$

15 wherein n_a is the number of amino acid alterations, x_a is the total number of amino acids in SEQ ID NO:2, y is 0.50 for 50%, 0.60 for 60%, 0.70 for 70%, 0.80 for 80%, 0.85 for 85%, 0.90 for 90%, 0.95 for 95%, 0.97 for 97% or 1.00 for 100%, and \bullet is the symbol for the multiplication operator, and wherein any non-integer product of x_a and y is rounded down to the nearest integer prior to subtracting it from x_a .

20 By way of example, a polypeptide sequence of the present invention may be identical to the reference sequence of SEQ ID NO:2, that is it may be 100% identical, or it may include up to a certain integer number of amino acid alterations as compared to the reference sequence such that the percent identity is less than 100% identity. Such
 25 alterations are selected from the group consisting of at least one amino acid deletion, substitution, including conservative and non-conservative substitution, or insertion, and wherein said alterations may occur at the amino- or carboxy-terminal positions of the reference polypeptide sequence or anywhere between those terminal positions, interspersed either individually among the amino acids in the reference sequence or in one

EXAMPLES:

The examples below are carried out using standard techniques, which are well known and
 5 routine to those of skill in the art, except where otherwise described in detail. The examples
 are illustrative, but do not limit the invention.

**Example 1: Sequence of BASB055 in *N. meningitidis* serogroup B strain
 ATCC13090.**

10 The sequence of the BASB055 gene of *N. meningitidis* strain ATCC13090 is shown in
 SEQ ID NO:1. The translation of the BASB055 polynucleotide sequence, showed in
 SEQ ID NO:2, is related by amino acid sequence homology to *Neisseria gonorrhoeae*
 MtrC protein. The BASB055 polypeptide contains a signal sequence characteristic of a
 lipoprotein.

Example 2: Construction of Plasmid to Express Recombinant BASB055

A: Cloning of BASB055.

The *Nde*I and *Xho*I restriction sites engineered into the forward Lip11-Fm/p (5'- AGG
 CAG AGG CAT ATG GCT TTT TAT GCT TTT AAG GCG ATG CG -3') ([SEQ ID
 20 NO:3]) and reverse Lip11-RCf/p (5'- AGG CAG AGG CTC GAG TTC CGC TTC
 AGA AGC AGT TTT GGC TTC -3')([SEQ ID NO:4]) amplification primers,
 respectively, permitted directional cloning of a BASB055 PCR product into the low
 copy number *E. coli* expression plasmid pTLZ2 such that a BASB055 protein could be
 expressed as a fusion protein containing a (His)₆ affinity chromatography tag at the C-
 25 terminus. The BASB055 PCR product was purified from the amplification reaction
 using silica gel-based spin columns (QiaGen) according to the manufacturers
 instructions. To produce the required *Nde*I and *Xho*I termini necessary for cloning,
 purified PCR product was sequentially digested to completion with *Nde*I and *Xho*I
 restriction enzymes as recommended by the manufacturer (Life Technologies).

Following the first restriction digestion, the PCR product was purified via spin column as above to remove salts and eluted in sterile water prior to the second enzyme digestion. The digested DNA fragment was again purified using silica gel-based spin columns prior to ligation with the pTLZ2 plasmid.

5

B: Production of Expression Vector.

To prepare the expression plasmid pTLZ2 for ligation, it was similarly digested to completion with both *NdeI* and *XhoI* and then treated with calf intestinal phosphatase (CIP, ~0.02 units / pmole of 5' end, Life Technologies) as directed by the manufacturer to prevent self-ligation. An approximately 5-fold molar excess of the digested fragment to the prepared vector was used to program the ligation reaction. A standard ~20 µl ligation reaction (~16°C, ~16 hours), using methods well known in the art, was performed using T4 DNA ligase (~2.0 units / reaction, Life Technologies). An aliquot of the ligation (~5 µl) was used to transform electro-competent BL21 DE3 cells according to methods well known in the art. Following a ~2-3 hour outgrowth period at 37°C in ~1.0 ml of LB broth, transformed cells were plated on LB agar plates containing ampicilline (100 µg/ml). The antibiotic was included in the selection media to ensure that all transformed cells carried the pTLZ2 plasmid (ApR). Plates were incubated overnight at 37°C for ~16 hours. Individual ApR colonies were picked with sterile toothpicks and used to "patch" inoculate fresh LB ApR plates as well as a ~1.0 ml LB ApR broth culture. Both the patch plates and the broth culture were incubated overnight at 37°C in either a standard incubator (plates) or a shaking water bath.

A whole cell-based PCR analysis was employed to verify that transformants contained the BASB055 DNA insert. Here, the ~1.0 ml overnight LB Ap broth culture was transferred to a 1.5 ml polypropylene tube and the cells collected by centrifugation in a Beckman microcentrifuge (~3 min., room temperature, ~12,000 X g). The cell pellet was suspended in ~200µl of sterile water and a ~10µl aliquot used to program a ~50µl final volume PCR reaction containing both BASB055 forward and reverse amplification

primers. Final concentrations of the PCR reaction components were essentially the same as those specified in example 2 except ~5.0 units of *Taq* polymerase was used. The initial 95°C denaturation step was increased to 3 minutes to ensure thermal disruption of the bacterial cells and liberation of plasmid DNA. An ABI Model 9700 thermal cycler and a 32 cycle, three-step thermal amplification profile, i.e. 95°C, 45sec; 55-58°C, 45sec, 72°C, 1min., were used to amplify the BASB055 PCR fragment from the lysed transformant samples. Following thermal amplification, a ~20µl aliquot of the reaction was analyzed by agarose gel electrophoresis (0.8 % agarose in a Tris-acetate-EDTA (TAE) buffer). DNA fragments were visualized by UV illumination after gel electrophoresis and ethidium bromide staining. A DNA molecular size standard (1 Kb ladder, Life Technologies) was electrophoresed in parallel with the test samples and was used to estimate the size of the PCR products. Transformants that produced the expected PCR product were identified as strains containing a BASB055 expression construct. Expression plasmid containing strains were then analyzed for the inducible expression of recombinant BASB055.

C: Expression Analysis of PCR-Positive Transformants.

For each PCR-positive transformant identified above, ~5.0 ml of LB broth containing ampicilline (100 µg/ml) was inoculated with cells from the patch plate and grown overnight at 37 °C with shaking (~250 rpm). An aliquot of the overnight seed culture (~1.0 ml) was inoculated into a 125 ml erlenmeyer flask containing ~25 ml of LB Ap broth and grown at 37 °C with shaking (~250 rpm) until the culture turbidity reached O.D.600 of ~0.5, i.e. mid-log phase (usually about 1.5 - 2.0 hours). At this time approximately half of the culture (~12.5 ml) was transferred to a second 125 ml flask and expression of recombinant BASB055 protein induced by the addition of IPTG (1.0 M stock prepared in sterile water, Sigma) to a final concentration of 1.0 mM. Incubation of both the IPTG-induced and non-induced cultures continued for an additional ~4 hours at 37 °C with shaking. Samples (~1.0 ml) of both induced and non-induced cultures were removed after the induction period and the cells collected by

centrifugation in a microcentrifuge at room temperature for ~3 minutes. Individual cell pellets were suspended in ~50µl of sterile water, then mixed with an equal volume of 2X Laemmli SDS-PAGE sample buffer containing 2-mercaptoethanol, and placed in boiling water bath for ~3 min to denature protein. Equal volumes (~15µl) of both the crude IPTG-induced and the non-induced cell lysates were loaded onto duplicate 12% Tris/glycine polyacrylamide gel (1 mm thick Mini-gels, Novex). The induced and non-induced lysate samples were electrophoresed together with prestained molecular weight markers (SeeBlue, Novex) under conventional conditions using a standard SDS/Tris/glycine running buffer (BioRad). Following electrophoresis, one gel was stained with commassie brilliant blue R250 (BioRad) and then destained to visualize novel BASB055 IPTG-inducible protein(s). The second gel was electroblotted onto a PVDF membrane (0.45 micron pore size, Novex) for ~2 hrs at 4 °C using a BioRad Mini-Protean II blotting apparatus and Towbin's methanol (20 %) transfer buffer. Blocking of the membrane and antibody incubations were performed according to methods well known in the art. A monoclonal anti- (His)5 antibody, followed by a second rabbit anti-mouse antibody conjugated to HRP (QiaGen), was used to confirm the expression and identity of the BASB055 recombinant protein. Visualization of the anti-His antibody reactive pattern was achieved using either an ABT insoluble substrate or using Hyperfilm with the Amersham ECL chemiluminescence system.

Example 3: Analysis of Recombinant BASB055 lipidation status

To determine if the BASB055 recombinant protein is lipidated, radiolabelling experiments using ³H-glycerol and ³H-palmitic acid as lipid-specific incorporation tag have been initiated. Cultures of the BL21 DE3 *E. coli* strain containing the expression plasmid was grown on M9 minimal medium and pulse-labelled with either 50µCi of ³H-glycerol or ³H-palmitic acid during the 3 hours of IPTG induction. Following a wash step to remove unincorporated isotope, whole cell lysates of radiolabelled cultures were run on gradient polyacrylamide gels (4-20%), fixed and then dried under vacuum. An autoradiograph was produced by exposing the dried gel to Hyperfilm (Amersham) at –

70°C for ~ about 7 days. A visible band of the same size as the IPTG-inducible coomassie stainable protein was detected by both labelling. That ³H-label incorporated into the IPTG inducible protein support the conclusion that the BASB055 protein is lipidated to some extent in *E. coli*.

5

Example 4: Production of Recombinant BASB055

Bacterial strain

- 10 A recombinant expression strain of *E. coli* BL21 DE3 containing a pTLZ2 plasmid encoding BASB055 from *N. meningitidis*. was used to produce cell mass for purification of recombinant protein. The expression strain was cultivated on LB agar plates containing 100µg/ml ampicilline ("Ap") to ensure plasmid maintenance. For cryopreservation at -80 °C, the strain was propagated in LB broth containing the same
- 15 concentration of antibiotic then mixed with an equal volume of LB broth containing 30% (w/v) glycerol.

Media

- The fermentation medium used for the production of recombinant protein consisted of
- 20 2X YT broth (Difco) containing 100µg/ml Ap. Antifoam was added to medium for the fermentor at 0.25 ml/L (Antifoam 204, Sigma). To induce expression of the BASB055 recombinant protein, IPTG (Isopropyl β-D-Thiogalactopyranoside) was added to the fermentor (1 mM, final).

Fermentation

- 25 A 500-ml erlenmeyer seed flask, containing 50ml working volume, was inoculated with 0.3 ml of rapidly thawed frozen culture, or several colonies from a selective agar plate culture, and incubated for approximately 12 hours at 37 ± 1°C on a shaking platform at 150rpm (Innova 2100, New Brunswick Scientific). This seed culture was then used to
- 30 inoculate a 5-L working volume fermentor containing 2X YT broth and both Ap

from Whatman Inc., Clifton, New Jersey. Dialysis tubing 8,000 MWCO was obtained from BioDesign Inc. Od New York, Carmal New York. BCA Protein Assay Reagents and Snake Skin dialysis tubing 3,500 MWCO were obtained from Pierce Chemical Co., Rockford, Illinois.

5

Extraction Protocol

Cell paste was thawed at room temperature for 30 to 60 minutes. Five to six grams of material was weighed out into a 50-ml disposable centrifuge tube. Recombinant BASB055 antigen was purified by extraction of cell membranes with 1.0 % Triton X114, and allowing phase partitioning based on Triton X114 cloud point at 37°C. The Triton X114 phase was diluted with 50 mM Tris-HCl containing 10% glycerol, 5% ethylene glycol and 0.5% Triton X100. This was applied to nickel-chelating Sepharose Fast Flow. The protein is afterwards eluted with 200 mM imidazole to affinity purify the histidine-tagged protein and yielded greater than 90% pure protein.

15

Final Formulation

BASB055 was formulated by dialysis overnight against, three changes of 0.1 % Triton X-100 and 1x PBS, pH 7.4. The purified protein was characterized and used to produce antibodies as described below.

20

Biochemical Characterizations : SDS-PAGE and Western Blot Analysis

The recombinant purified protein was resolved on 4-20 % polyacrylamide gels and electrophoretically transferred to PVDF membranes at 100 V for 1 hour as previously described (Thebaine et al. 1979, Proc. Natl. Acad. Sci. USA 76:4350-4354). The PVDF membranes were then pretreated with 25 ml of Dulbecco's phosphate buffered saline containing 5 % non-fat dry milk. All subsequent incubations were carried out using this pretreatment buffer.

25

PVDF membranes were incubated with a dilution of anti-His tail antibodies for 1 hour at room temperature. PVDF membranes were then washed twice with wash buffer (20 mM Tris buffer, pH 7.5, containing 150 mM sodium chloride and 0.05 % Tween-20). PVDF membranes were incubated with 25 ml of a 1:5000 dilution of peroxidase-labeled species specific conjugate for 30 minutes at room temperature. PVDF membranes were then washed 4 times with wash buffer, and were developed with 3-amino-9-ethylcarbazole and urea peroxide as supplied by Zymed (San Francisco, CA) for 10 minutes each.

The results of an SDS-PAGE (Figure 1A) show a protein about 50 kDa purified to greater than 90 % and that is reactive to an anti-tetra (His) antibody by western blots (Figure 1B) of the SDS-PAGE.

Example 5 : Immunization of mice with recombinant BASB055

Partially purified recombinant BASB055 protein expressed in *E. coli* has been injected three times in Balb/C mice on days 0, 14 and 28 (10 animals/group). Animals were injected by the subcutaneous route with around 5 µg of antigen in two different formulations: either adsorbed on 100 µg AlPO₄ or formulated in SBAS2 emulsion (SB62 emulsion containing 5 µg MPL and 5µg QS21 per dose). A negative control group consisting of mice immunized with the SBAS2 emulsion only has also been added in the experiment. Mice were bled on days 28 (14 days Post II) and 35 (7 days Post III) in order to detect specific anti-BASB055 antibodies. Specific anti-BASB055 antibodies were measured by Elisa on partially purified BASB055 protein as well as on *E. coli* proteins. Antibody responses were also evaluated by western-blotting when tested against different *Neisseria meningitidis* B strains. Pooled sera (from 10 mice/group) from both formulations (on day 7 Post III only) were tested in these assays. Results illustrated in Figure 2 clearly indicate that the antibody response is quite good, while the anti-*E. coli* antibody response (Figure 3), which is clearly positive, is much lower than the specific BASB055 response (Figure 2). The AlPO₄ formulation induces

WO 00/43517

PCT/EP00/00425

the highest antibody levels. Western-blotting illustrated in Figures 4 and 5 confirm that the BASB055 protein is well recognized at the expected MW of 50 kDa by immunized mice sera.

Recognition of the BASB055 epitopes on different *Neisseria meningitidis* serogroup B strains by western-blotting

In this test, immunized mice sera (pooled) have been tested by western-blotting for recognition of the BASB055 epitopes on seven different *Neisseria meningitidis* B strains: H44/76 (B:15:P1.7, 16, lineage ET-5), M97 250687 (B:4:P1.15), BZ10 (B:2b:P1.2, lineage A4), BZ198 (B:NT*: -, lineage 3), EG328 (B:NT*, lineage ST-18), NGP165 (B:2a:P1.2, ET 37 cluster) and the ATCC 13090 (B:15:P1.15) *Neisseria meningitidis* B strains, as well as on partially purified recombinant BASB055 protein. (* : NT : Not Typed).

Briefly, 10 µl (> 10⁸ cells/lane) of each sample treated with sample buffer (10 min at 95°C) are put into a SDS-PAGE gradient gel (Tris-glycine 4-20%, Novex, code n°EC6028). Electrophoretic migration occurs at 125 volts for 90 min. (35mA/gel). Afterwards, proteins are transferred to nitrocellulose sheet (0.45 µm, Bio-rad code n° 162-0114) at 100 volts for 1:30 hour using a Bio-rad Trans-blot system (code n°170-3930). Filter was blocked with PBS - 0.05 % Tween 20 overnight at room temperature, before incubation with the mice sera containing the anti-BASB055 antibodies from both ALPO4 and SBAS2 formulations. These sera are diluted 100 times in PBS - 0.05 % Tween 20, and incubated on the nitrocellulose sheet for two hours at room temperature with gentle shaking. After three repeated washing steps in PBS - 0.05 % Tween 20 for 5 min., the nitrocellulose sheet is incubated at room temperature for 1 hour under gentle shaking with the appropriate conjugate (biotinylated anti-mouse Ig antibodies from sheep, Amersham code n°RPN1001) diluted at 1/500 in the same washing buffer. The membrane is washed three times as previously, and incubated for 30 min with agitation using the streptavidin-peroxidase complex (Amersham code n°1051) diluted at 1/1000 in the washing buffer. After the last three repeated washing steps, the revelation occurs during the 20 min incubation time in a 50 ml solution containing 30 mg 4-chloro-1-naphthol (Sigma), 10 ml methanol, 40 ml PBS, and 30 µl of H₂O₂. The staining is stopped while washing the membrane several times in distilled water.

Results illustrated in Figures 4 and 5 show that all strains tested present a clear reactivity against the major BASB055 band around 50 kDa, plus few other bands at lower MW, which are most probably related to the antigen, as they are not present on the *E. coli* preparation (Figure 5). These results suggest that the BASB055 protein is expressed in probably all *Neisseria meningitidis* B strains. In both Figures 4 and 5, the recombinant BASB055 protein is recognized by mice sera at the same MW. However, the lowest band recognized by mice sera (< 9 kDa) appears to be related to an *E. coli* contaminant, as demonstrated in the last lane (Figure 5) with *E. coli* extract.

WO 00/43517

PCT/EP00/00425

SEQUENCE INFORMATION

BASB055 Polynucleotide and Polypeptide Sequences

SEQ ID NO:1

5 *Neisseria meningitidis* BASB055 polynucleotide sequence from strain ATCC 13090

ATGGCTTTTTATGCTTTTAAGGCGATGCGTGCGGCCGCGTTGGCTGCCGCCGTTGCATTG
 GTACTGTCTGCTCTTGCGGTAAAGGCGGAGACGCGGCGCAGGGCGGGCAGCCTGCTGGTCCG
 GAAGCCCCTGCGCCCCGTCGTCGGTGTCTGTAACCGTCCATCCGCAAACCGTCGCATTGACC
 GTCGAGTTGCCGGGGCGTTTGAATCGCTGCGTACCGCCGATGTCCGCGCCCAAGTCGGC
 10 GGCATCATCCAAAACGCCTGTTCCAAGAAGGCAGTTATGTCCGTGCCGGACAGCCGCTG
 TATCAGATCGACAGTTCCACTTATGAAGCAAATCTGGAAAGCGCGCGCGCAACTGGCA
 ACGGCTCAGGCAACGCTTGCCAAAGCGGATGCGGATTGGCGCGATACAAGCCTTTGGTT
 GCCGCCGAAGCGTCAGCCGGCAGGAATACGATGCTGCGGTAACGGCGAAACGTTCTGCC
 GAGGCAGGTGTCAAAGCAGCACAGGCGGCAATCAAATCTGCCGGCATTAACTGAACCGT
 15 TCGCGCATTACCGCGCGGATTTCCGGCTTTATCGGTCACTCCAAAGTTTCCGAAGGTACG
 CTGTTGAATGCGGGCGATACGACCGTGCTGGCAACCATCCGCCAAACCAATCCGATGTAT
 GTGAACGTTACCCAGTCTGCATCCGAAGTGATGAAATTGCGCCGTGAGATAGCCGAAGGC
 AAAGTCTGCGCGCGGATGGTGTGATTGCGGTGCGCATCAAATTTGACGACGGCACAGTT
 TACCCTGAAAAAGGCCGCTGCTGTTTGCCGATCCGGTCGTCAACGAATCGACCGGTCAG
 20 ATTACCCTGCGCGCCCGCTACCGAACGATCAGAATATCCTGATGCCCGGTCTGTATGTG
 CGCGTGCTGATGGACCAAGTGGCGGTGGATAACGCATTTGTTGTGCCGAGCAGGCGGTA
 ACGCGCGGTGCGAAAGATACCGTGATGATTGTGAATGCCCAAGGCGGTATGGAACCCCGC
 GAGGTAACGGTTGCGCAACAGCAGGATACGAATTGGATTGTTACGTGCGGTCTGAAGGAC
 GGGGACAAGGTGGTGTGGAAGGCATCAGTATCGCCGGTATAACGGGTGCGAAAAAGGTA
 25 ACGCCCAAAGAAATGGGCGTCTGTGAAAACCAAGCCCGCGCCTCAATCCGGCGTTTCA
 ACGGCATCTGAAGCCAAAACCTGCTTCTGAAGCGGAATAA

SEQ ID NO:2

Neisseria meningitidis BASB055 polypeptide sequence deduced from the polynucleotide of

30 SeQ ID NO:1

MAFYAFKAMRAAALAAVALVLSSCGKGGDAAQGGQPAGREAPAPVVGVTVPQTVALT
 VELPGRLESLRTADVRAQVGIIQKRLFQEGSYVRAGQPLYQIDSSTYEANLESARAQLA
 TAQATLAKADADLARYKPLVAEAVSRQYDAAVTAKRSAEAGVKAAQAAIKSAGINLNR
 35 SRITAPISGFIGQSKVSEGTLNAGDTTVLATIRQTNPMYVNVTSASEVMKLRRQIAEG
 KLLAADGVIAVGIFKDDGTVPYPEKGRLLFADPVVNESTGQITLRAAVPNDQNILMPGLYV
 RVLMDQVAVDNAFVVPQAVTRGAKDTVMIVNAQGGMEPREVTVAQQQGTNWIIVTSGLKD
 GDKVVVEGISIAGITGAKKVTPEKAWASSENQAAAPQSGVQTASEAKTASEAE

40 SEQ ID NO:3

AGG CAG AGG CAT ATG GCT TTT TAT GCT TTT AAG GCG ATG CG

SEQ ID NO:4

45 AGG CAG AGG CTC GAG TTC CGC TTC AGA AGC AGT TTT GGC TTC

A deposit containing a *Neisseria meningitidis* Serogroup B strain has been deposited with the American Type Culture Collection (herein "ATCC") on June 22, 1997 and assigned deposit number 13090. The deposit was described as *Neisseria meningitidis* (Albrecht and Ghon) and is a freeze-dried, 1.5-2.9 kb insert library constructed from *N. meningitidis* isolate. The deposit is described in Int. Bull. Bacteriol. Nomencl. Taxon. 8: 1-15 (1958).

10

15

20

INDICATIONS RELATING TO DEPOSITED MICROORGANISM
OR OTHER BIOLOGICAL MATERIAL

(PCT Rule 13bis)

A. The indications made below relate to the deposited microorganism or other biological material referred to in the description on page <u>64</u> , line <u>2-21</u>	
B. IDENTIFICATION OF DEPOSIT Further deposits are identified on an additional sheet <input type="checkbox"/>	
Name of depositary institution AMERICAN TYPE CULTURE COLLECTION	
Address of depositary institution (including postal code and country) 10801 UNIVERSITY BLVD, MANASSAS, VIRGINIA 20110-2209, UNITED STATES OF AMERICA	
Date of deposit 22 June 1997 (22.06.97)	Accession Number 13090
C. ADDITIONAL INDICATIONS (leave blank if not applicable) This information is continued on an additional sheet <input type="checkbox"/>	
In respect of those designations where a European Patent is sought, a sample of the deposited microorganism will be made available until the publication of the mention of the grant of the European Patent or until the date on which the application has been refused or withdrawn, only by issue of such a sample to an expert nominated by the person requesting the sample.	
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States)	
E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable)	
The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g., "Accession Number of Deposit")	

For receiving Office use only
<input type="checkbox"/> This sheet was received with the international application
Authorized officer

For International Bureau use only
<input type="checkbox"/> This sheet was received by the International Bureau on:
Authorized officer

CLAIMS

1. An isolated polypeptide comprising an amino acid sequence which has at least 85% identity to the amino acid sequence of: SEQ ID NO:2.
- 5 2. An isolated polypeptide as claimed in claim 1 in which the amino acid sequence has at least 95% identity to the amino acid of: SEQ ID NO:2.
3. The polypeptide as claimed in claim 1 comprising the amino acid of: SEQ ID NO: 2.
- 10 4. An isolated polypeptide of SEQ ID NO:2.
5. An immunogenic fragment of the polypeptide as claimed in any one of claims 1 to 4 in which the immunogenic activity of said immunogenic fragment is substantially the same as the polypeptide of SEQ ID NO:2.
- 15 6. An isolated polynucleotide comprising a nucleotide sequence encoding the polypeptide of SEQ ID NO:2.
- 20 7. An isolated polynucleotide comprising the polynucleotide of SEQ ID NO:1.
8. An isolated polynucleotide comprising a nucleotide sequence encoding the polypeptide of SEQ ID NO:2, obtainable by screening an appropriate library under stringent hybridization conditions with a labeled probe having the sequence of SEQ ID NO:1 or a fragment thereof.
- 25 9. An expression vector or a recombinant live microorganism comprising an isolated polynucleotide according to any one of claims 6 - 8.

10. A process for expressing a polynucleotide of any one of claims 6 - 8 comprising transforming a host cell with the expression vector comprising at least one of said polynucleotides and culturing said host cell under conditions sufficient for expression of
5 any one of said polynucleotides.
11. A vaccine composition comprising an effective amount of the polypeptide of any one of claims 1 to 5 and a pharmaceutically acceptable carrier.
- 10 12. A vaccine composition comprising an effective amount of the polynucleotide of any one of claims 6 to 8 and a pharmaceutically acceptable carrier.
13. The vaccine composition according to either one of claims 11 or 12 wherein said composition comprises at least one other *Neisseria meningitidis* antigen.
15
14. An antibody immunospecific for the polypeptide or immunological fragment as claimed in any one of claims 1 to 5.
15. A method of diagnosing a *Neisseria meningitidis* infection, comprising identifying a
20 polypeptide as claimed in any one of claims 1 - 5, or an antibody that is immunospecific for said polypeptide, present within a biological sample from an animal suspected of having such an infection.
16. Use of a composition comprising an immunologically effective amount of a
25 polypeptide as claimed in any one of claims 1 - 5 in the preparation of a medicament for use in generating an immune response in an animal.

17. Use of a composition comprising an immunologically effective amount of a polynucleotide as claimed in any one of claims 6 - 8 in the preparation of a medicament for use in generating an immune response in an animal.
- 5 18. A therapeutic composition useful in treating humans with *Neisseria meningitidis* disease comprising at least one antibody directed against the polypeptide of claims 1 - 5 and a suitable pharmaceutical carrier.

PCT


WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau




INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : C12N 15/31, C07K 14/22, 16/12, G01N 33/68, A61K 39/095, 39/40, 31/70		A1	(11) International Publication Number: WO 00/43517
			(43) International Publication Date: 27 July 2000 (27.07.00)
(21) International Application Number: PCT/EP00/00425		(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).	
(22) International Filing Date: 19 January 2000 (19.01.00)			
(30) Priority Data: 9901462.3 22 January 1999 (22.01.99) GB 9902069.5 29 January 1999 (29.01.99) GB			
(71) Applicant (for all designated States except US): SMITHKLINE BEECHAM BIOLOGICALS S.A. [BE/BE]; Rue de l'Institut 89, B-1330 Rixensart (BE).			
(72) Inventor; and (75) Inventor/Applicant (for US only): THONNARD, Joelle [BE/BE]; SmithKline Beecham Biologicals s.a., Rue de L'Institut 89, B-1330 Rixensart (BE).			
(74) Agent: PRIVETT, Kathryn, Louise; SmithKline Beecham, Two New Horizons Court, Brentford, Middlesex TW8 9EP (GB).		Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>	
(54) Title: BASB055 POLYNUCLEOTIDE AND POLYPEPTIDE FROM NEISSERIA MENINGITIDIS. USES THEREOF			
(57) Abstract The invention provides BASB055 polypeptides and polynucleotides encoding BASB055 polypeptides and methods for producing such polypeptides by recombinant techniques. Also provided are diagnostic, prophylactic and therapeutic uses.			

Figure 1: Analysis of recombinant, purified BASB055. A. Coomassie stained SDS-polyacrylamide gel of purified BASB055. B. Western blot of purified BASB055 with anti histidine immune reagent.



09/889756

Figure 2 : Anti-BASB055 antibodies induced in Balb/C mice

09/889756


Figure 3 : Anti-*E. coli* response induced in BASB055 immunized mice sera

Figure 4 : Recognition of the BASB055 protein on several *Neisseria meningitidis* strains with BASB055 immunized mice sera.





Figure 5 : Recognition of the BASB055 protein on several *Neisseria meningitidis* strains with BASB055 immunized mice sera.



Docket No.: BM45353

PCT/EP00/00425

DECLARATION AND POWER OF ATTORNEY

As a below named inventor, I hereby declare that:

My residence, post office address and citizenship are as stated below next to my name.

I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled:

BASB055 POLYNUCLEOTIDE AND POLYPEPTIDE FROM NEISSERIA MENINGITIDIS.
USES THEREOF

the specification of which (check one)

☐ is attached hereto.

☒ was filed on 19 January 2000 as Serial No. PCT/EP00/00425
and was amended on (if applicable).

I hereby state that I have reviewed and understand the contents of the above identified specification, including the claims, as amended by any amendment referred to above.

I acknowledge the duty to disclose information which is material to the patentability as defined in Title 37, Code of Federal Regulations, Section 1.56.

I hereby claim foreign priority benefits under Title 35, United States Code, Section 119(a)-(d) or Section 365(b) of any foreign application(s) for patent or inventor's certificate, or Section 365(a) of any PCT International application which designated at least one country other than the United States, listed below and have also identified below any foreign application for patent or inventor's certificate, or PCT International application having a filing date before that of the application on which priority is claimed.

Prior Foreign Application(s)

Number	Country	Filing Date	Priority Claimed
9901462.3	Great Britain	22 January 1999	Yes
9902069.5	Great Britain	29 January 1999	Yes

I hereby claim the benefit under Title 35, United States Code, Section 119(e) of any United States provisional application(s) listed below.

Application Number	Filing Date
--------------------	-------------

I hereby claim the benefit under Title 35, United States Code, Section 120 of any United States application(s) or Section 365(c) of any PCT International application designating the United States, listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States or PCT International application in the manner provided by the

first paragraph of Title 35, United States Code, Section 112, I acknowledge the duty to disclose information which is material to patentability as defined in Title 37, Code of Federal Regulations, Section 1.56 which became available between the filing date of the prior application and the national or PCT international filing date of this application.


Serial No. Filing Date Status

I hereby appoint the practitioners associated with the Customer Number provided below to prosecute this application and to transact all business in the Patent and Trademark Office connected therewith, and direct that all correspondence be addressed to that Customer Number:

Customer Number 25,308

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

1-a
Full Name of Inventor: Joelle THONNARD

Inventor's Signature: 

Date: 10-07-2007

Residence: Rixensart, Belgium

BEX

Citizenship: Belgian

Post Office Address: SmithKline Beecham Corporation
Corporate Intellectual Property - UW2220
P.O. Box 1539
King of Prussia, Pennsylvania 19406-0939

SEQUENCE LISTING

<110> SmithKline Beecham Biologicals S.A.

<120> Novel compounds

<130> BM45353

<160> 4

<170> FastSEQ for Windows Version 3.0

<210> 1

<211> 1239

<212> DNA

<213> Neisseria meningitidis

<400> 1

atggcttttt atgcttttaa ggcgatgcgt gcggccgcgt tggctgccgc cgttgcatcg	60
gtactgtcgt cttgcggtaa aggcggagac gcggcgagg gcgggcagcc tgctggtcgg	120
gaagccccctg cgcgcgtcgt cgggtgtcgt accgtccatc cgcaaaccgt cgcattgacc	180
gtcgagttgc cggggcgctt ggaatcgctg cgtaccgcgc atgtccgcgc ccaagtcggc	240
ggcatcatcc aaaaacgcct gttccaagaa ggcagttatg tccgtgccgg acagccgctg	300
tatcagatcg acagttccac ttatgaagca aatctggaaa gcgcgcgcgc gcaactggca	360
acggctcagg caacgcttgc caaagcggat gcggatttgg cgcgatacaa gcctttggtt	420
gccgccgaag ccgtcagccg gcaggaatac gatgctgcgg taacggcgaa acgttctgcc	480
gaggcaggtg tcaaagcagc acaggcggca atcaaatctg ccggcattaa tctgaaccgt	540
tcgcgcatta ccgcgcgat ttccggcttt atcggtcagt ccaaagtctc cgaaggtacg	600
ctgttgaaatg cgggcgatac gaccgtgctg gcaaccatcc gccaaaccaa tccgatgtat	660
gtgaacgtta ccagtcctgc atccgaagtg atgaaattgc gccgtcagat agccgaaggc	720
aaactgctgg cggcggatgg tgtgattgcg gtcggcatca aatttgacga cggcacagtt	780
taccctgaaa aaggccgcct gctgtttgcc gatccggctg tcaacgaatc gaccggtcag	840
attaccctgc gcgcgcgcgt accgaacgat cagaatatcc tgatgcccg tctgtatgtg	900
cgcgtgctga tggaccaagt ggcggtggat aacgcatttg ttgtgccga gcaggcggta	960
acgcgcggtg cgaaagatac cgtgatgatt gtgaatgcc aaggcggat ggaacccgc	1020
gaggtaacgg ttgcgcaaca gcagggtacg aattggattg ttacgtcggg tctgaaggac	1080
ggggacaagg tggttggtga aggcatacgt atcgcggta taacgggtgc gaaaaaggta	1140
acgccccaaag aatgggcgtc gtctgaaaac caagccgcgc cgcctcaatc cggcgttcag	1200

